

New U.S. Non-Provisional Utility Patent Application

Title: **METHODS FOR REGULATING DIFFERENTIATION OF NEURAL
CELLS AND USES THEREOF**

Inventors: Lloyd A. Greene and James M. Angelastro

METHODS FOR REGULATING DIFFERENTIATION OF NEURAL CELLS AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Serial Number 60/460,242 filed

5 April 4, 2003.

STATEMENT OF GOVERNMENT INTEREST

[0001] This invention was made with government support under NIH/NINCDS Grant No. NS-16036. As such, the United States government has certain rights in this invention.

BACKGROUND OF THE INVENTION

10 [0002] A key step in the formation of the nervous system is the determination of proliferating neural progenitor cells to undergo differentiation into neurons and glia. Despite major advances in identification and characterization of neural progenitor cells (Placzek and Furley, Patterning cascades in the neural tube. *Neural development. Curr. Biol.* 6:526-29, 1996; Gage, F.H., Mammalian neural stem cells. *Science*, 287:1433-38, 2000; Kintner, C., 15 Neurogenesis in embryos and in adult neural stem cells. *J. Neurosci.*, 22:639-43, 2002; Schuurmans and Guillemot, Molecular mechanisms underlying cell fate specification in the developing telencephalon. *Curr. Opin. Neurobiol.*, 12:26-34, 2002), the mechanisms that govern this determination are only partially understood.

[0003] The selective degeneration of specific types or classes of neurons of the 20 central nervous system (CNS) underlies many neurological disorders. This realization has generated interest in defining populations of progenitor cells that, through manipulation of the neuronal differentiation process, may serve as replenishable sources of neurons, and, therefore, may present an option for treating neurodegenerative and demyelinating disorders. Additionally, it is well recognized that neural tumors and other cancers develop when cells 25 divide and grow uncontrollably. Thus, a means of manipulating the differentiation process of tumor cells may provide a therapy for the treatment of cancers.

[0004] Neural degeneration may result from neurodegenerative diseases, CNS 30 traumas, and the acquired secondary effects of non-neural dysfunction. Alzheimer's disease is a neurodegenerative disease characterized by a progressive, inexorable loss of cognitive function. The pathogenesis of Alzheimer's disease is associated with an excessive number of

neuritic, or senile, plaques (composed of neurites, astrocytes, and glial cells around an amyloid core) in the cerebral cortex, and neurofibrillary tangles (composed of paired helical filaments). Approximately 4 million Americans suffer from Alzheimer's disease, at an annual cost of about \$90 billion. The disease is about twice as common in women as in men, and accounts for more than 65% of the dementias in the elderly. While senile plaques and neurofibrillary tangles occur with normal aging, they are much more prevalent in persons with Alzheimer's disease. To date, a cure for Alzheimer's disease is not available, and cognitive decline is inevitable.

[0005] Demyelination is also a feature of many neurologic disorders. Demyelinating conditions are manifested in loss of myelin – the multiple dense layers of lipids and protein which cover many nerve fibers. Multiple sclerosis (MS) is the most prevalent demyelinating condition. In Europe and North America, an average of 40-100 people out of every 100,000 have MS. The disease affects approximately 250,000 people in the United States alone. Histopathologically, MS is characterized by inflammation, plaques of demyelination infiltrating cells in the CNS tissue, loss of oligodendroglia, and focal axonal injury. Typically, the symptoms of MS include lack of co-ordination, paresthesias, speech and visual disturbances, and weakness. Current treatments for the various demyelinating conditions are often expensive, symptomatic, and only partially effective, and may cause undesirable secondary effects. Corticosteroids represent the main form of therapy for MS. While these may shorten the symptomatic period during attacks, they may not affect eventual long-term disability. Long-term corticosteroid treatment is rarely justified, and can cause numerous medical complications, including osteoporosis, ulcers, and diabetes.

[0006] This year, and each year in the foreseeable future, 17,000 people in the United States will be diagnosed with brain tumors. Brain tumors invade and destroy normal tissue, producing such effects as impaired sensorimotor and cognitive function, increased intracranial pressure, cerebral edema, and compression of brain tissue, cranial nerves, and cerebral vessels. Drowsiness, lethargy, obtuseness, personality changes, disordered conduct, and impaired mental faculties are the initial symptoms in 25% of patients with malignant brain tumors. Treatment of brain tumors is often multimodal, and depends on pathology and location of the tumors. For malignant gliomas, multimodal therapy, including chemotherapy, radiation therapy, and surgery, is used to try to reduce tumor mass. Regardless of approach,

however, prognosis for patients suffering from these tumors is guarded: the median term of survival after chemotherapy, radiation therapy, and surgery is only about 1 year, and only 25% of these patients survive for 2 years.

[0007] In view of the foregoing, it is clear that many neural disorders are related to loss of cells, loss of myelin, or loss of cell control. An ability to regulate the differentiation of neuroprogenitor cells into various differentiated neural cells would provide supplies of neural cells that could be effective in treating such neural disorders. However, prior to the present invention, the direction of such progenitor cells along specific pathways of neuronal differentiation has proved difficult.

SUMMARY OF THE INVENTION

[0008] The inventors disclose herein that the b-zip transcription factor, ATF5, plays a major regulatory role in the differentiation of neuroprogenitor cells into differentiated neural cells. In particular, the inventors have discovered that, in the developing brain, ATF5 expression is high within ventricular zones containing neural stem cells and neural progenitor cells, but is undetectable in post-mitotic neurons. In attached clonal neurosphere cultures, ATF5 is expressed by neural stem cells and neural progenitor cells, but is undetectable in tau-positive neurons. In PC12 cell cultures, nerve growth factor (NGF) dramatically down-regulates endogenous ATF5 protein and transcripts, while exogenous ATF5 suppresses NGF-promoted neurite outgrowth. Such inhibition may require repression of cyclic AMP (cAMP) responsive element (CRE) DNA-binding sites and/or other ATF5 DNA-binding sites, including those not yet discovered. By contrast, loss of function conferred by dominant-negative ATF5 accelerates NGF-promoted neuritogenesis. Exogenous ATF5 suppresses neurogenesis by cultured nestin-positive telencephalic cells, while dominant-negative ATF5, and a small interfering RNA targeted to ATF5, promote this activity. These findings indicate that ATF5 blocks differentiation of neuroprogenitor cells into neurons, and must be down-regulated to permit this process to occur. Additional studies carried out in culture, and also *in vivo*, indicate that ATF5 blocks differentiation of neural progenitor cells into differentiated astroglia and oligodendroglia, and that dominant-negative ATF5 accelerates this differentiation.

[0009] Accordingly, the present invention provides a method for promoting differentiation of a neural stem cell or a neural progenitor cell into a differentiated neural cell,

by inhibiting ATF5 in the cell. Also provided is a differentiated neural cell produced by this method.

[0010] The present invention also provides a method for producing differentiated neural cells by: (a) obtaining or generating a culture of neural stem cells or neural progenitor cells; (b) contacting the culture of neural stem cells or neural progenitor cells with an amount of an ATF5 inhibitor effective to produce differentiated neural cells; and (c) optionally, contacting the differentiated neural cells with at least one neurotrophic factor. Examples of methods for contacting the cells with (treating the cells with) the ATF5 inhibitor or the neurotrophic factor (in protein or nucleic acid form) include, without limitation, absorption, electroporation, immersion, injection, liposome delivery, transfection, vectors, and other protein-delivery and nucleic-acid-delivery vehicles and methods. Also provided is a population of cells, comprising the differentiated neural cells produced by this method.

[0011] The present invention further provides a method for treating nervous tissue degeneration in a subject in need of treatment by: (a) obtaining or generating a culture of neural stem cells or neural progenitor cells; (b) contacting the culture of neural stem cells or neural progenitor cells with an amount of an ATF5 inhibitor effective to produce differentiated neural cells; (c) optionally, contacting the differentiated neural cells with at least one neurotrophic factor; and (d) transplanting the differentiated neural cells into the subject in an amount effective to treat the nervous tissue degeneration.

[0012] Additionally, the present invention provides differentiated neural cells produced by: (a) obtaining or generating a culture of neural stem cells or neural progenitor cells; (b) contacting the neural stem cells or neural progenitor cells with an amount of an ATF5 inhibitor effective to produce differentiated neural cells; and (c) optionally, contacting the differentiated neural cells with at least one neurotrophic factor. Also provided are a transgenic non-human animal containing these differentiated neural cells, and uses of these differentiated neural cells in analyzing neuron development, function, and death, and in monitoring synaptic differentiation.

[0013] The present invention is also directed to a method for isolating and/or purifying a population of differentiated neural cells by: (a) obtaining or generating a culture of neural stem cells or neural progenitor cells that express enhanced green fluorescent protein (eGFP); (b) contacting the culture of neural stem cells or neural progenitor cells with an

amount of an ATF5 inhibitor effective to produce differentiated neural cells that express eGFP; (c) optionally, contacting the differentiated neural cells with at least one neurotrophic factor; (d) detecting expression of eGFP in the differentiated neural cells; and (e) isolating the differentiated neural cells that express eGFP.

5 [0014] Furthermore, the present invention provides a method for identifying an agent for use in treating a condition associated with nervous tissue degeneration by: (a) obtaining or generating a culture of neural stem cells or neural progenitor cells; (b) contacting the neural stem cells or neural progenitor cells with an amount of an ATF5 inhibitor effective to produce neurons, wherein some or all of the neurons are degenerated; (c) contacting the
10 degenerated neurons with a candidate agent; and (d) determining if the agent enhances regeneration or survival of some or all of the degenerated neurons.

[0015] The present invention also provides a method for suppressing differentiation of neural stem cells or neural progenitor cells into differentiated neural cells, by contacting the neural stem cells or neural progenitor cells with an amount of ATF5 effective to suppress
15 differentiation in the neural stem cells or neural progenitor cells.

[0016] Additionally, the present invention is directed to a therapeutic composition, comprising: (a) a nucleic acid encoding an ATF5 inhibitor; (b) a vector; and (c) optionally, a pharmaceutically-acceptable carrier. Also provided is a method for treating a neural tumor in a subject in need of treatment, by administering the therapeutic composition to the subject.

20 [0017] The present invention further provides a method for identifying an agent which inhibits ATF5 by: (a) contacting a candidate agent with ATF5, in the presence of CRE; and (b) assessing the ability of the candidate agent to inhibit interaction between ATF5 and CRE. This method may further comprise the steps of: (c) contacting the candidate agent with neural stem cells or neural progenitor cells containing ATF5; and (d) determining if the agent
25 has an effect on an ATF5-associated biological event in the cells. Also provided are agents identified by these methods, as well as methods for promoting differentiation in neural stem cells or neural progenitor cells, and for treating or preventing a neural tumor in a subject, using these agents.

[0018] Furthermore, the present invention provides a method for determining whether
30 a subject has a neural tumor, by assaying a diagnostic sample of the subject for ATF5, wherein detection of an ATF5 level elevated above normal is diagnostic of a neural tumor in

the subject. Also provided are methods for assessing the efficacy of therapy to treat a neural tumor in a subject who has undergone or is undergoing treatment for a neural tumor, and for assessing the prognosis of a subject who has a neural tumor.

[0019] Finally, the present invention provides a kit for use in detecting a neural tumor, comprising: (a) an ATF5-specific agent; and (b) reagents suitable for detecting ATF5; wherein the ATF5-specific agent is selected from the group consisting of an agent reactive with ATF5 and a nucleic acid probe which hybridizes to nucleic acid encoding ATF5.

[0020] Additional aspects of the present invention will be apparent in view of the description which follows.

BRIEF DESCRIPTION OF THE FIGURES

[0021] FIG. 1 shows that nerve growth factor (NGF) down-regulates ATF5 protein in PC12 cells, and demonstrates a reciprocal relationship with neurite outgrowth. (A) Time course of the effect of NGF treatment on ATF5 protein expression in PC12 cells. Cells were exposed to NGF for the indicated times, and 135 μ g of whole cell extracts were subjected to Western immunoblotting, first with anti-ATF5, then, after stripping, with anti-ERK to normalize for loading. Numbers at the left of the figure indicate the positions of molecular weight markers (in kDa). Comparable results were achieved in 3 independent experiments. (B) Comparison of the kinetics of NGF-dependent down-regulation of ATF5 expression, and promotion of neurite outgrowth. The relative levels of ATF5 expression were determined by densitometry, and normalized to levels of ERK protein in the same sample; the levels are reported in arbitrary units. Proportions of cells bearing neurites of a length at least twice the diameter of the cell body were determined in the same cultures, by scoring at least 200 cells per time point.

[0022] FIG. 2 shows that overexpression of ATF5 represses neurite outgrowth in PC12 cells, while NTAzip-ATF5 accelerates neuritogenesis. (A) Detection and NGF response of PC12 cells expressing exogenous ATF5. PC12 cells were transiently transfected with pCMS-*eGFP* (panels a and b) or pCMS-*eGFP* expressing FLAG-tagged ATF5 (panels c and d). Two days after transfection, the cultures were treated with NGF. Five days after transfection (*i.e.*, after 3 days of NGF exposure), the cells were fixed and co-stained with rabbit anti-GFP (panels a and c) or mouse anti-FLAG antibody (panels b and d), with detection by FITC (GFP) and rhodamine-conjugated secondary antibody (FLAG-ATF5).

-7-

scale bar represents 50 μm (B) Quantification of the effects of exogenous ATF5 and of NTAzip-ATF5 on NGF-promoted neurite outgrowth. PC12 cells were transiently transfected with pCMS-eGFP, without insert or expressing FLAG-tagged ATF5 or FLAG-tagged NTAzip-ATF5. Two days after transfection, the cultures were treated with NGF. Cultures were fixed at the indicated times, after commencement of NGF exposure, and immunostained with anti-GFP and anti-FLAG, as above. Transfected cells (positive for FLAG and/or GFP staining) were assessed for the presence or absence of neurites. The proportions of transfected cells bearing neurites are reported \pm SEM, with $n = 3$ cultures (and at least 300 transfected cells assessed per culture). Comparable results were achieved in 4 additional independent experiments. In all cases (including the data shown), ANOVA analysis indicated a p value of <0.05 at the 72-h point of NGF treatment for eGFP vs. ATF5. (C) NTAzip accelerates NGF-promoted neurite outgrowth. Cultures were transfected, treated, and assessed as in (B), at 24 h after NGF exposure. Values represent the mean \pm SEM for results of 4 independent experiments. In each experiment, the data were normalized to the percentage of neurite-bearing cells transfected with pCMS-eGFP. The average percentage of such cells was 10.6 ± 3.7 . NTAzip vs. eGFP: $p < 0.02$, Student's t-distribution test

[0023] FIG. 3 illustrates that ATF5 is differentially expressed in the ventricular zones of E12-E15 rat brain. (A) Expression of ATF5 message in developing rat brain (panels a and b). *In situ* hybridization was carried out using an ATF5 antisense probe in sagittal sections of E15 rat brain. Panel a shows the area around the fourth ventricle, and panel b shows the telencephalon. There was no positive signal with a control ATF5 sense probe. Expression of ATF5 protein is shown in coronal sections of E12 (panels c and d) and E14 (panels e and f) rat telencephalon. (panel c) Staining with pre-immune serum. (panels d-f) Co-staining with anti-ATF5 (red) and anti-tubulin β (III) (TUJ1 antibody; green). Arrows indicate staining of ATF5 in the ventricular zone (VZ). CX = cortex; scale bar for panel a represents 100 μm (B) High-power confocal images of reciprocal expression of ATF5 (red) and tubulin β (III) in coronal sections of E14 rat telencephalon. Immunohistochemical staining was carried out as in (A). Images showing the ventricular zone (panel a) and cortex (panel b) are from the same section, and were photographed in the same confocal Z-plane section (1.3 μm). Arrowhead shows a migratory cell undergoing a transition from a progenitor to a neuron, by exhibiting both ATF5 and tubulin β (III) staining. Co-localization was confirmed by YZ and XZ confocal images. scale bar for panel B represents 20 μm

[0024] FIG. 4 demonstrates reciprocal expression of ATF5 and tubulin β (III) in E17 rat brain. (A-C) Expression of ATF5 (red) and tubulin β (III) (green) in the area of the anterior (A-C) and posterior (D-F) lateral ventricles of the E17 rat brain.

Immunohistochemical staining was carried out as in FIG. 3 and the Examples. scale bar
5 represents 100 μ m

[0025] FIG. 5 shows that ATF5 is expressed in neural stem cells and progenitor cells, but not in mature neurons in attached neurosphere cultures. Attached clonal neurosphere cultures were established from the subventricular zone and hippocampal dentate gyrus of newborn mouse brain, and maintained as described in the Examples. Cultures were fixed and
10 co-stained as follows: (A) ATF5 (red) and AC133 (green), a stem cell marker. Thick arrows show examples of nuclear staining, thin arrows show cytoplasmic staining. (B) ATF5 (red) and nestin (green), a marker for neural progenitor cells. Arrows indicate nuclear staining. (C, D) ATF5 (red) and NF-M (green), a marker for the neuronal lineage. Arrows show nuclear staining in (C) and cell body in (D). (E, F) ATF5 (red) and anti-tau (green), a
15 neuronal marker. Comparable results were achieved in 10 independent experiments. Arrows show neurons at the periphery of the cultures; arrowhead shows stem and neural progenitor cells at the center of the culture. Stained cells were examined and photographed by confocal microscopy. The scale bar is 20 μ m for (A), and 50 μ m for (B-F).

[0026] FIG. 6 illustrates that ATF5 represses, and NTAzip-ATF5 promotes, neurite
20 outgrowth and expression of neuronal markers in neural progenitor cells. (A) Cultured E14 telencephalic cells were transiently transfected with pCMS-*eGFP* containing either no insert (empty vector), *FLAG-ATF5*, or *NTAzip-ATF5*. Three days following transfection, the cultures were fixed and co-immunostained for GFP and either nestin or tubulin β (III) (TUJ1 antibody). Transfected cells (GFP+) were assessed for the presence of neurite-like processes,
25 and for co-expression of the indicated markers. Values represent the mean \pm SEM for 3 cultures in which at least 300 transfected cells were evaluated per culture. Comparable results were achieved in 4 independent experiments. ANOVA analysis of transfected cells: total cells – nestin/*eGFP* alone vs. nestin/*ATF5*, $p < 0.001$; TUJ1/*eGFP* alone vs. TUJ1/*ATF5*, $p < 0.05$; nestin/*eGFP* alone vs. nestin/*NTAzip*, and TUJ1/*eGFP* alone vs. TUJ1/*NTAzip*, no
30 significant difference; process-bearing cells – TUJ1/*GFP* alone vs. TUJ1/*ATF5*, $p < 0.05$; nestin/*eGFP* alone vs. nestin/*NTAzip* and TUJ1/*eGFP* alone vs. TUJ1/*NTAzip*, no significant

difference. (B) Cultured E14 telencephalic cells were infected with retroviruses expressing eGFP or FLAG-ATF5 and eGFP. One week after infection, the cultures were fixed and assessed as in (A), and assessed for NF-M expression. Comparable results were achieved in 3 independent experiments. ANOVA analysis: total cells – nestin/eGFP alone vs. nestin/ATF5, $p < 0.001$; TUJ1/GFP alone vs. TUJ1/ATF5, $p < 0.01$; NFM/eGFP vs. NFM/ATF5, $p < 0.001$; process-bearing cells – TUJ1/GFP alone vs. TUJ1/ATF5, $p < 0.001$; NFM/GFP alone vs. NFM/ATF5, $p < 0.01$; nestin/eGFP alone vs. nestin/ATF5, no significant difference; TUJ1 vs. NFM, no significance both with eGFP alone and with eGFP plus ATF5. (C) E14 telencephalon cells were infected with retroviruses expressing eGFP, eGFP and FLAG-ATF5, or eGFP-FLAG-NTAzip-ATF5. Four days after infection, the cultures were fixed and evaluated as in (A). Comparable results were achieved in 2 independent experiments. ANOVA analysis: nestin/GFP alone vs. nestin/ATF5, $p < 0.001$; total and process-bearing cells – TUJ1/eGFP alone vs. TUJ1/ATF5, $p < 0.01$; TUJ1/GFP alone vs. TUJ1/NTAzip, $p < 0.05$. (D) Cultured E14 telencephalic cells were transiently transfected with pCMS-eGFP, with or without ATF5 siRNA. Four days following transfection, the cultures were fixed and co-immunostained either for GFP and TUJ1 antibody, or with GFP and ATF5 antiserum. Transfected cells (GFP+) were assessed for the presence of the neuronal marker, tubulin β (III) (TUJ1), or ATF5. Values represent the mean \pm SEM for six cultures in which at least 300 transfected cells were evaluated per culture. Comparable results were achieved in 3 independent experiments (two experiments with E14 telencephalon cells cultured with serum plus EGF and FGF2, and one experiment with only serum). ANOVA analysis: TUJ1/eGFP alone vs. TUJ1/ATF5 siRNA, $p < 0.001$; ATF5/GFP alone vs. ATF5/ATF5 siRNA, $p < 0.001$. (E) ATF5 suppresses NT3-promoted neuronal differentiation. E15 telencephalon cells were infected with retroviruses expressing eGFP, eGFP and FLAG-ATF5, or eGFP-FLAG-NTAzip-ATF5, all \pm NT3. Four days after infection and maintenance \pm NT3 treatment, the cultures were fixed and evaluated, as in (A), for eGFP and TUJ1 expression. Comparable results were achieved in 2 independent experiments. ANOVA analysis: -NT3/eGFP alone vs. +NT3/GFP alone, $p < 0.001$; -NT3/eGFP alone vs. -NT3/ATF5, $p < 0.05$; +NT3/eGFP alone vs. +NT3/ATF5, $p < 0.001$; -NT3/eGFP alone vs. -NT3/NTAzip, $p < 0.001$; +NT3/GFP alone vs. +NT3/NTAzip, no significant difference.

[0027] FIG. 7 demonstrates that NTAzip-ATF5 and VP16-CREB reverse ATF5-promoted repression of CRE-mediated gene expression and of neurite outgrowth. (A) PC12

cells were co-transfected with pGI3-CRE *luciferase*, pcDNA-*LacZ*, and 1 μ g / culture of pCMS-*eGFP* expressing either no insert (empty vector), *FLAG-ATF5*, *FLAG-NTAzip-ATF5*, or *VP16-CREB*. The cultures were also exposed to NGF for 2 days prior to and during the time of transfection (for a total of 3 days), during the time of transfection (1 day), or during the last hour prior to harvesting. One day after transfection, the cells were harvested and assessed for luciferase expression and *LacZ* activity (β -GAL). Values represent mean normalized CRE-luciferase activity (in arbitrary units) \pm SEM (n = 3). Comparable results were achieved in three independent experiments. Student's t-distribution test: empty vector (eGFP alone) vs. VP16CREB at all times, $p < 0.001$; GFP alone vs. ATF5, $p < 0.033$ by day 3.

(B) PC12 cells were co-transfected with pGI3-CRE *luciferase*, pcDNA-*LacZ*, and the indicated combinations of pCMS-*eGFP* expressing either no insert (GFP), *FLAG-ATF5* (*ATF5*), *FLAG-NTAzip-ATF5* (*AZIP*), or *VP16-CREB*. The latter vectors were each used at 0.5 μ g / culture, and empty vector was added, as needed, to bring the total DNA level to 1 μ g / culture. Cultures were harvested 1 day later, or assayed for luciferase expression and *LacZ* activity (β -GAL). Where indicated, NGF was added to the medium 1 h before harvesting. Values represent mean normalized CRE-luciferase activity (in arbitrary units) \pm SEM (n = 6), with data pooled from 2 independent experiments. Student's t-distribution test: -NGF - eGFP alone vs. ATF5, $p < 0.003$; eGFP alone vs. NTAzip, $p < 0.0003$; eGFP alone vs. ATF5/NTAzip, no significant difference; eGFP alone vs. VP16CREB and VP16CREB/ATF5, $p < 0.0001$; +NGF - eGFP alone vs. ATF5, $p < 0.0001$; eGFP alone vs. NTAzip, $p < 0.02$; eGFP alone vs. ATF5/NTAzip, $p < 0.02$; eGFP alone vs. VP16CREB and VP16CREB/ATF5, $p < 0.0001$. (C) PC12 cells were co-transfected with the indicated constructs, and NGF was added to the medium 2 days later. Transfected cells (identified for eGFP) were assessed for neurite outgrowth at the indicated times. Values represent means \pm SEM of results for 3 cultures in which at least 300 transfected cells were scored per culture. Comparable results were obtained in 2 independent experiments. ANOVA analysis after 72 h of NGF-treatment: eGFP alone vs. ATF5, eGFP $p < 0.001$; eGFP alone vs. NTAzip-ATF5, NTAzip/ATF5, Vp16CREB, or Vp16CREB/ATF5, no significant difference.

[0028] FIG. 8 sets forth the nucleotide sequence of ATF5.

[0029] FIG. 9 sets forth the amino acid sequence of ATF5.

DETAILED DESCRIPTION OF THE INVENTION

[0030] As described above, a key step in the formation of the nervous system is the determination of proliferating neural progenitor cells to exit the cell cycle and undergo neuronal differentiation. Despite major advances in identification and characterization of such progenitor cells, the mechanisms that govern this determination are only partially understood.

[0031] One system with potential to address this issue is the PC12 line of pheochromocytoma cells (Greene and Tischler, Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor. *Proc. Natl. Acad. Sci. U S A*, 73:2424-28, 1976; Burstein and Greene, Evidence for RNA synthesis-dependent and -independent pathways in stimulation of neurite outgrowth by nerve growth factor. *Proc. Natl. Acad. Sci. U S A*, 75:6059-63, 1978). In the presence of the neurotrophic factor, nerve growth factor (NGF), proliferating neuroblast-like PC12 cells acquire, by means of a transcription-dependent mechanism, a neuronal phenotype characterized by formation of axons, up-regulation of a number of neuronal markers, and transition to a post-mitotic state.

[0032] To identify genes responsible for this neuronal differentiation, the inventors employed serial analysis of gene expression (SAGE) to provide a comprehensive profile and comparison of transcripts present in PC12 cells, before and after 9 days of treatment with NGF (Angelastro *et al.*, Identification of diverse nerve growth factor-regulated genes by serial analysis of gene expression (SAGE) profiling. *Proc. Natl. Acad. Sci. U S A*, 97:10424-29, 2000). Of the approximately 22,000 unique transcripts detected in the cells, approximately 4% underwent a 6-fold or greater increase or decrease in expression after NGF exposure. Among the identified genes with the greatest change in expression was ATF5, a member of the activating transcription factor (ATF/CREB) family. In response to NGF, ATF5 transcripts, which were among the most highly expressed in the cells prior to treatment, fell by 25-fold in relative expression.

[0033] Relatively few studies have been carried out to characterize ATF5 (also known as ATFX and ATF-7) and its biological functions (Nishizawa and Nagata, cDNA clones encoding leucine-zipper proteins which interact with G-CSF gene promoter element 1-binding protein. *FEBS Lett.*, 299:36-38, 1992; Pati *et al.*, Human Cdc34 and Rad6B ubiquitin-conjugating enzymes target repressors of cyclic AMP-induced transcription for

proteolysis. *Mol. Cell Biol.*, 19:5001-13, 1999; Peters *et al.*, ATF-7, a novel bZIP protein, interacts with the PRL-1 protein-tyrosine phosphatase. *J. Biol. Chem.*, 276:13718-726, 2001; Persengiev *et al.*, Inhibition of apoptosis by ATFx: a novel role for a member of the ATF/CREB family of mammalian bZIP transcription factors. *Genes Dev.*, 16:1806-14, 2002).

5 ATF5 is a b-zip transcription factor that forms homodimers that, at least *in vitro*, bind cyclic AMP (cAMP) responsive element (CRE) DNA-binding sites. In addition, ATF5 represses cAMP-induced transcription in intact cells (Pati *et al.*, Human Cdc34 and Rad6B ubiquitin-conjugating enzymes target repressors of cyclic AMP-induced transcription for proteolysis. *Mol. Cell Biol.*, 19:5001-13, 1999; Peters *et al.*, ATF-7, a novel bZIP protein, interacts with
10 the PRL-1 protein-tyrosine phosphatase. *J. Biol. Chem.*, 276:13718-726, 2001), and has been shown to inhibit apoptosis (Persengiev *et al.*, Inhibition of apoptosis by ATFx: a novel role for a member of the ATF/CREB family of mammalian bZIP transcription factors. *Genes Dev.*, 16:1806-14, 2002). This raised the possibility that ATF5 might interfere with the activity of transcription factors, such as CREB, that appear to promote neuronal
15 differentiation *via* CRE-mediated gene activation (Finkbeiner *et al.*, CREB: a major mediator of neuronal neurotrophin responses. *Neuron*, 19:1031-47, 1997; Dawson and Ginty, CREB family transcription factors inhibit neuronal suicide. *Nat. Med.*, 8:450-51, 2002; Lonze *et al.*, Apoptosis, axonal growth defects, and degeneration of peripheral neurons in mice lacking CREB. *Neuron*, 34:371-85, 2002). These properties, along with its down-regulation by NGF,
20 suggest that ATF5 is a negative regulator of neuronal differentiation, *via* CRE and other (as yet undiscovered) DNA-binding sites.

[0034] The present invention relates to several findings concerning the levels of expression of ATF5 in cells of the nervous system. In particular, the inventors have discovered that ATF5 is highly expressed in the nuclei of neuroprogenitor cells (in both the
25 developing and adult nervous systems), and that it functions in these cells to block their differentiation into neurons, astroglia, and oligodendroglia. In contrast, ATF5 is only detected outside the nucleus in oligodendroglia and Schwann cells (myelin-forming cells in the CNS and the peripheral nervous system (PNS), respectively), and is not detected in mature neurons or astroglia. Studies also indicate that ATF5 is highly expressed in human
30 neuroblastoma cells.

[0035] The present invention also relates to a role for ATF5 in the differentiation of neuroprogenitor cells. For example, the inventors have observed that forced constitutive expression of ATF5 protein in neuroprogenitor cells blocks their differentiation into neurons and glial cells. The inventors have also observed that specific suppression of ATF5 protein synthesis, or forced constitutive expression of a blocking form of the protein, strongly promotes differentiation of neuroprogenitor cells.

[0036] Furthermore, the present invention relates to regulation of ATF5 expression. In particular, the inventors' findings indicate that ATF5 expression is regulated by neurotrophic factors, and, therefore, is an essential part of the mechanism by which they promote neuronal differentiation.

[0037] Accordingly, the present invention provides a method for promoting differentiation of a neural stem cell or a neural progenitor cell into a differentiated neural cell, as well as a differentiated neural cell produced by this method. Differentiation is the cellular process by which cells become structurally and functionally specialized during development. As used herein, the term "promoting differentiation" means activating, enhancing, inducing, initiating, or stimulating differentiation of a neural stem cell or a neural progenitor cell.

[0038] Neural stem cells are cultured cells, derived from the pluripotent inner cell mass of blastocyst stage embryos, that are capable of replicating indefinitely. In general, neural cells have the potential to differentiate into neural cells (*i.e.*, they are pluripotent); thus, they may serve as a continuous source of new neural cells. The neural stem cell of the present invention may be obtained from any animal, but is preferably obtained from a mammal (*e.g.*, human, domestic animal, or commercial animal). In one embodiment of the present invention, the neural stem cell is a murine neural stem cell. In another, preferred, embodiment, the neural stem cell is obtained from a human.

[0039] As used herein, a "differentiated neural cell" is a partially-differentiated or fully-differentiated cell of the central nervous system (CNS) or peripheral nervous system (PNS), and includes, without limitation, a fully-differentiated ganglion cell, glial (or neuroglial) cell (*e.g.*, an astrocyte, astroglial cell, oligodendrocyte, oligodendroglial cell, or Schwann cell), granule cell, neuronal cell (or neuron), and stellate cell, as well as any neural progenitor cells thereof. Progenitor cells are parent cells which, during development and differentiation, give rise to a distinct cell lineage by a series of cell divisions. Neural

progenitor cells, for example, are committed to a cell lineage that will develop, eventually, into fully-differentiated neural cells of the CNS or PNS; however, such neural progenitor cells may not yet be dedicated to a particular type, or subclass, of neural cell.

[0040] Initially, neural progenitor cells may acquire a rostral character (*e.g.*, rostral
5 neural progenitor cells), followed by a positional identity (*e.g.*, cerebellar progenitor cells, cerebral progenitor cells, or spinal progenitor cells). Such partially-differentiated neural progenitor cells may become committed to a cell line that will differentiate into a specific type of neural cell (*e.g.*, progenitor cells of astrocytes, astroglial cells, ganglion cells, granule cells, neurons, oligodendrocytes, oligodendroglial cells, Schwann cells, or stellate cells), and,
10 thereafter, give rise to fully-differentiated neural cells (*e.g.*, astrocytes, astroglial cells, ganglion cells, granule cells, neurons, oligodendrocytes, oligodendroglial cells, Schwann cells, or stellate cells). Accordingly, the partially-differentiated neural cell of the present invention may be a cell, with a neural identity, that has acquired a directional or positional character, or that has committed to developing into a particular class of neural cell, but is not
15 a fully-differentiated neural cell.

[0041] The neural progenitor cell of the present invention may be obtained from any animal, but is preferably obtained from a mammal (*e.g.*, human, domestic animal, or commercial animal). In one embodiment of the present invention, the neural progenitor cell is a murine neural progenitor cell. In another, preferred, embodiment, the neural progenitor
20 cell is obtained from a human.

[0042] A "neuronal cell", or "neuron", as used herein, is a conducting or nerve cell of the nervous system that typically consists of a cell body (perikaryon) that contains the nucleus and surrounding cytoplasm; several short, radiating processes (dendrites); and one long process (the axon), which terminates in twig-like branches (telodendrons), and which
25 may have branches (collaterals) projecting along its course. Examples of neurons include, without limitation, cerebellar neurons, or neurons from the cerebellum (*e.g.*, basket cells, Golgi cells, granule cells, Purkinje cells, and stellate cells); cortical neurons, or neurons from the cerebral cortex (*e.g.*, pyramidal cells and stellate cells, including interneurons, midbrain neurons, and neurons of the substantia nigra); hippocampal cells, or cells from the
30 hippocampus (including granule cells); cells of the Pons; neurons of the dorsal root ganglia (DRG); motor neurons; peripheral neurons; sensory neurons; neurons of the spinal cord;

ventral interneurons; and primary neurons (neurons taken directly from the brain, and, in general, placed into a tissue culture dish), all of which may be cholinergic, dopaminergic, GABAergic, or serotonergic.

[0043] Differentiation of neural stem cells and neural progenitor cells into partially-
5 or fully-differentiated neural cells may be detected by known cellular or molecular
procedures, and assays and methods disclosed herein. In one embodiment of the present
invention, the differentiated neural cell is a post-mitotic neuron. The term "post-mitotic", as
used herein, refers to a neuron that is in G0 phase (a quiescent state), and is no longer
dividing or cycling. In another embodiment of the present invention, the differentiated neural
10 cell is genetically marked, in that it expresses enhanced green fluorescent protein (eGFP), as
described herein. The eGFP genetic marker may be particularly useful in a method for
isolating and/or purifying a population of differentiated neural cells, as described below.

[0044] The method of the present invention comprises inhibiting ATF5 in a neural
stem cell or neural progenitor cell. As used herein, "ATF5" includes both an "ATF5 protein"
15 and an "ATF5 analogue". Unless otherwise indicated, "protein" shall include a protein,
protein domain, polypeptide, or peptide, and any fragment thereof. The ATF5 protein has the
amino acid sequence set forth in FIG. 9, including conservative substitutions thereof. As
described below, Western immunoblotting permitted the inventors to deduce the major
cellular form of ATF5 protein. The ATF5 cDNA sequence predicts two potential in-frame
20 methionine start sites that would lead to proteins of approximately 30 and 20 kDa. The
inventors' observation that the major form of ATF5 in cells has an apparent molecular mass
of 20-22 kDa indicates favored utilization of the second site. When a canonical Kozak
initiation consensus sequence was included upstream of the first methionine, the larger
protein was expressed, thereby indicating that the 22-kDa form is not formed by cleavage of a
25 30-kDa precursor. Accordingly, the ATF5 protein of the present invention further includes
both the 22-kDa and 30-kDa isomers thereof.

[0045] As used herein, "conservative substitutions" are those amino acid substitutions
which are functionally equivalent to a substituted amino acid residue, either because they
have similar polarity or steric arrangement, or because they belong to the same class as the
30 substituted residue (*e.g.*, hydrophobic, acidic, or basic). The term "conservative
substitutions" includes substitutions having an inconsequential effect on the ability of ATF5

to interact with CRE, particularly in respect of the use of said interaction for the identification and design of agonists of ATF5, for molecular replacement analyses, and/or for homology modeling.

[0046] An "ATF5 analogue", as used herein, is a functional variant of the ATF5 protein, having ATF5 biological activity, that has 60% or greater (preferably, 70% or greater) amino-acid-sequence homology with the ATF5 protein. As further used herein, the term "ATF5 biological activity" refers to the activity of a protein or peptide that demonstrates an ability to associate physically with, or bind with, CRE (*i.e.*, binding of approximately two fold, or, more preferably, approximately five fold, above the background binding of a negative control), under the conditions of the assays described herein, although affinity may be different from that of ATF5.

[0047] It will be obvious to the skilled practitioner that the numbering of amino acid residues in ATF5, or in the ATF5 analogues or peptidomimetics covered by the present invention, may be different than that set forth herein, or may contain certain conservative amino acid substitutions that produce the same ATF5-CRE associating activity as that described herein. Corresponding amino acids and conservative substitutions in other isoforms or analogues are easily identified by visually inspecting the relevant amino acid sequences, or by using commercially available homology software programs.

[0048] In accordance with methods described herein, ATF5 may be inhibited in a neural stem cell or neural progenitor cell by disabling, disrupting, or inactivating the function or activity of ATF5 in the cell, or by diminishing the amount or level of ATF5 in the cell. For example, ATF5 in a cell may be inhibited by targeting ATF5 directly. Additionally, activity of ATF5 in a cell may be inhibited indirectly, by targeting an enzyme or other endogenous molecule that regulates or modulates the functions or levels of ATF5 in the cell. ATF5 expression may also be inhibited by engineering the ATF5 gene so that ATF5 is expressed on an inducible promoter. In such a case, ATF5 expression would be sustained in the presence of a suitable inducing agent, but would shut down once the supply of inducer was depleted, thereby resulting in a decrease in the amount or level of ATF5 in the cell.

[0049] Preferably, activity of the ATF5 in the neuron is inhibited or decreased by at least 10% in the method of the present invention. More preferably, activity of the ATF5 is decreased by at least 20%. Activity of the ATF5 is inhibited in the neural stem cell or neural

progenitor cell by an amount effective to promote differentiation of the neural stem cell or neural progenitor cell. This amount may be readily determined by the skilled artisan, based upon known procedures, including analysis of titration curves established *in vivo*, and methods disclosed herein.

5 [0050] By way of example, activity of the ATF5 in a neuron may be inhibited by directly or indirectly inactivating, interfering with, or down-regulating the CRE-binding function of ATF5 in the neural stem cell or neural progenitor cell (*e.g.*, by the modulation or regulation of proteins that interact with ATF5). The ATF5 in a neural stem cell or neural progenitor cell may be inactivated, for example, by contacting the neural stem cell or neural
10 progenitor cell with a small molecule or protein mimetic that inhibits ATF5 or that is reactive with (*i.e.*, has affinity for, binds to, or is directed against) ATF5. Examples of methods for contacting the cell with (treating the cell with) a molecule or protein mimetic include, without limitation, absorption, electroporation, immersion, injection, liposome delivery, transfection, vectors, and other protein-delivery and nucleic-acid-delivery vehicles and methods, as
15 described below.

[0051] Activity of ATF5 in a neural stem cell or neural progenitor cell also may be inhibited by directly or indirectly causing, inducing, or stimulating the down-regulation of ATF5 expression within the cell. Accordingly, in one embodiment of the present invention, activity of ATF5 is inhibited in a neural stem cell or neural progenitor cell by contacting the
20 cell with a modulator of ATF5 expression, in an amount effective to promote differentiation of the cell. The modulator may be a protein, polypeptide, peptide, nucleic acid (including DNA or RNA), antibody, Fab fragment, F(ab')₂ fragment, molecule, compound, antibiotic, drug, or an agent reactive with (*i.e.*, has affinity for, binds to, or is directed against) ATF5, that inhibits or down-regulates ATF5 expression. A Fab fragment is a univalent antigen-
25 binding fragment of an antibody, which is produced by papain digestion. An F(ab')₂ fragment is a divalent antigen-binding fragment of an antibody, which is produced by pepsin digestion.

[0052] Modulators of ATF5 may be identified using a simple screening assay. For example, to screen for candidate modulators of ATF5, neural progenitor
30 cells may be plated onto microtiter plates, then contacted with a library of drugs. Any resulting decrease in, or down-regulation of, ATF5 expression then may be

-18-

detected using a luminescence reporter, nucleic acid hybridization, and/or immunological techniques known in the art, including an ELISA. Additional modulators of ATF5 expression may be identified using screening procedures well known in the art or disclosed herein. Modulators of ATF5 will include those
5 drugs which inhibit or down-regulate expression of ATF5. In this manner, candidate modulators also may be screened for their ability to promote differentiation of neural stem cells or neural progenitor cells, and, therefore, their ability to treat neural tumors, as discussed below.

[0053] In one embodiment of the present invention, ATF5 in a neural stem cell or
10 neural progenitor cell is inhibited by contacting the cell with an ATF5 inhibitor. As used herein, "an ATF5 inhibitor" shall include a protein, polypeptide, peptide, nucleic acid (including DNA, RNA, and an antisense oligonucleotide), antibody (monoclonal and polyclonal, as described above), Fab fragment (as described above), F(ab')₂ fragment (as described above), molecule, compound, antibiotic, drug, and any combinations thereof, and
15 may be an agent reactive with ATF5, as defined above. By way of example, the ATF5 inhibitor of the present invention may be a neurotrophic factor. As used herein, a "neurotrophic factor" is a factor involved in the nutrition or maintenance of neural tissue. Neurotrophic factors, may further the development and differentiation of committed neural progenitor cells, or they may induce or enhance the growth and survival of differentiated
20 neural cells. A classic example of a neurotrophic factor is NGF (nerve growth factor). Other examples of neurotrophic factors for use in the present invention include, without limitation, GDNF, NT3, CNTF, and BDNF, as well as cognate receptors thereof (including TrkB and TrkC). These factors may be obtained from R&D Systems, Inc. (Minneapolis, MN).

[0054] Additionally, the ATF5 inhibitor of the present invention may be an ATF5
25 transgene, comprising the ATF5 gene and an inducible promoter, in the absence of a suitable inducer. In a cell containing such a transgene, ATF5 expression would be sustained in the presence of a suitable inducing agent; however, ATF5 expression would be shut down once the supply of inducer was depleted. Thus, an ATF5 transgene, comprising the ATF5 gene and an inducible promoter, would, in the absence of a suitable inducer, effectively bring
30 about a decrease in the amount or level of ATF5 in the cell, thereby functioning as an ATF5 inhibitor.

[0055] The ATF5 inhibitor of the present invention also may be an interfering RNA, or RNAi, including ATF5 small interfering RNA (siRNA), as disclosed herein. As used herein, "RNAi" refers to a double-stranded RNA (dsRNA) duplex of any length, with or without single-strand overhangs, wherein at least one strand, putatively the antisense strand, is homologous to the target mRNA to be degraded. As further used herein, a "double-stranded RNA" molecule includes any RNA molecule, fragment, or segment containing two strands forming an RNA duplex, notwithstanding the presence of single-stranded overhangs of unpaired nucleotides. Additionally, as used herein, a double-stranded RNA molecule includes single-stranded RNA molecules forming functional stem-loop structures, such that they thereby form the structural equivalent of an RNA duplex with single-strand overhangs. The double-stranded RNA molecule of the present invention may be very large, comprising thousands of nucleotides; preferably, however, it is small, in the range of 21-25 nucleotides. In a preferred embodiment, the RNAi of the present invention comprises a double-stranded RNA duplex of at least 19 nucleotides.

[0056] In one embodiment of the present invention, RNAi is produced *in vivo* by an expression vector containing a gene-silencing cassette coding for RNAi (see, *e.g.*, U.S. Patent No. 6,278,039, *C. elegans* deletion mutants; U.S. Patent Application No. 2002/0006664, Arrayed transfection method and uses related thereto; WO 99/32619, Genetic inhibition by double-stranded RNA; WO 01/29058, RNA interference pathway genes as tools for targeted genetic interference; WO 01/68836, Methods and compositions for RNA interference; and WO 01/96584, Materials and methods for the control of nematodes). In another embodiment of the present invention, RNAi is produced *in vitro*, synthetically or recombinantly, and transferred into the microorganism using standard molecular-biology techniques. Methods of making and transferring RNAi are well known in the art. See, *e.g.*, Ashrafi *et al.*, Genome-wide RNAi analysis of *Caenorhabditis elegans* fat regulatory genes. *Nature*, 421:268-72, 2003; Cottrell *et al.*, Silence of the strands: RNA interference in eukaryotic pathogens. *Trends Microbiol.*, 11:37-43, 2003; Nikolaev *et al.*, Parc. A Cytoplasmic Anchor for p53. *Cell*, 112:29-40, 2003; Wilda *et al.*, Killing of leukemic cells with a BCR/ABL fusion gene RNA interference (RNAi). *Oncogene*, 21:5716-24, 2002; Escobar *et al.*, RNAi-mediated oncogene silencing confers resistance to crown gall tumorigenesis. *Proc. Natl. Acad. Sci. USA*, 98:13437-42, 2001; and Billy *et al.*, Specific interference with gene expression induced

by long, double-stranded RNA in mouse embryonal teratocarcinoma cell lines. *Proc. Natl. Acad. Sci. USA*, 98:14428-33, 2001.

[0057] Furthermore, the ATF5 inhibitor of the present invention may be an oligonucleotide antisense to ATF5. Oligonucleotides antisense to ATF5 may be designed
5 based on the nucleotide sequence of ATF5, which is readily available (FIG. 8). For example, a partial sequence of the ATF5 nucleotide sequence (generally, 18-20 base pairs), or a variation sequence thereof, may be selected for the design of an antisense oligonucleotide. This portion of the ATF5 nucleotide sequence may be within the 5' domain. A nucleotide sequence complementary to the selected partial sequence of the ATF5 gene, or the selected
10 variation sequence, then may be chemically synthesized using one of a variety of techniques known to those skilled in the art, including, without limitation, automated synthesis of oligonucleotides having sequences which correspond to a partial sequence of the ATF5 nucleotide sequence, or a variation sequence thereof, using commercially-available oligonucleotide synthesizers, such as the Applied Biosystems Model 392 DNA/RNA
15 synthesizer.

[0058] Once the desired antisense oligonucleotide has been prepared, its ability to inhibit ATF5 then may be assayed. For example, the oligonucleotide antisense to ATF5 may be contacted with neural progenitor cells, and the levels of ATF5 expression or activity in the cells may be determined using standard techniques, such as Western-blot analysis and
20 immunostaining. Alternatively, the antisense oligonucleotide may be delivered to neural progenitor cells using a liposome vehicle, then the levels of ATF5 expression or activity in the cells may be determined using standard techniques, such as Western-blot analysis. Where the level of ATF5 expression in the cells is reduced in the presence of the designed antisense oligonucleotide, it may be concluded that the oligonucleotide could be a useful ATF5
25 inhibitor.

[0059] It is within the confines of the present invention that oligonucleotide antisense to ATF5 may be linked to another agent, such as a drug or a ribozyme, in order to increase the effectiveness of treatments using ATF5 inhibition, increase the efficacy of targeting, and/or increase the efficacy of degradation of ATF5 RNA. Examples of antineoplastic drugs
30 to which the antisense oligonucleotide may be linked include, without limitation, carboplatin, cyclophosphamide, doxorubicin, etoposide, and vincristine. Moreover, oligonucleotide

antisense to ATF5 may be prepared using modified bases (*e.g.*, a phosphorothioate) to make the oligonucleotide more stable and better able to withstand degradation.

[0060] The ATF5 inhibitor of the present invention also may be a dominant-negative form of the protein (*e.g.*, NTAzip-ATF5), as disclosed herein. In one embodiment, the
5 dominant-negative form of ATF5 is expressed on an inducible promoter.

[0061] Additional ATF5 inhibitors may be identified using screening procedures well known in the art, and methods described below.

[0062] The present invention contemplates the use of proteins and protein analogues generated by synthesis of polypeptides *in vitro*, *e.g.*, by chemical means or *in vitro* translation
10 of mRNA. For example, ATF5 and inhibitors thereof may be synthesized by methods commonly known to one skilled in the art (*Modern Techniques of Peptide and Amino Acid Analysis* (New York: John Wiley & Sons, 1981); Bodansky, M., *Principles of Peptide Synthesis* (New York: Springer-Verlag New York, Inc., 1984). Examples of methods that may be employed in the synthesis of the amino acid sequences, and analogues
15 of these sequences, include, but are not limited to, solid-phase peptide synthesis, solution-method peptide synthesis, and synthesis using any of the commercially-available peptide synthesizers. The amino acid sequences of the present invention may contain coupling agents and protecting groups, which are used in the synthesis of protein sequences, and which are well known to one of skill in the art.

20 [0063] The method of the present invention comprises inhibiting ATF5 in a neural stem cell or neural progenitor cell by contacting the cell with an ATF5 inhibitor. The inhibitor is provided in an amount effective to produce a differentiated neural cell. This amount may be readily determined by the skilled artisan, based upon known procedures and methods disclosed herein. The inventors have demonstrated herein that neurons cultured in
25 the presence of neurotrophic factors survive and elaborate processes. Accordingly, in another embodiment, the method of the present invention further comprises the step of contacting the neural stem cell or neural progenitor cell with at least one neurotrophic factor, contemporaneously with, or following, inhibition of ATF5. The neurotrophic factors of the present invention are provided in amounts effective to produce a fully-differentiated neural
30 cell of the CNS or PNS (*e.g.*, a neuron). These amounts may be readily determined by the skilled artisan, based upon known procedures and methods disclosed herein.

[0064] In the method of the present invention, neural stem cells or neural progenitor cells may be contacted with effective amounts of an ATF5 inhibitor and neurotrophic factors *in vitro*, or *in vivo* in a subject. The inhibitor and factors may be contacted with a neural stem cell or neural progenitor cell by introducing the inhibitor and factors into the cell. Where
5 contacting is effected *in vitro*, the inhibitor and factors may be added directly to the culture medium, as described herein. Alternatively, the inhibitor and factors may be contacted with a neural stem cell or neural progenitor cell *in vivo* in a subject, by introducing the inhibitor and factors into the subject (*e.g.*, by introducing the inhibitor and factors into cells of the subject), or by administering the inhibitor
10 and factors to the subject. The subject may be any neural or developed animal, but is preferably a mammal (*e.g.*, a human, domestic animal, or commercial animal). More preferably, the subject is a human.

[0065] Where the inhibitor and factors are contacted with the cell *in vivo*, the subject is preferably an embryo. However, it is within the confines of the present invention for the
15 cells to be transplanted into a fully-grown human or animal subject, and for the inhibitor and factors then to be administered to the human in order to effect differentiation of the neural stem cells or neural progenitor cells into differentiated neural cells *in vivo* in the subject. The cells may be contained in nervous tissue of a subject, and may be detected in nervous tissue of the subject by standard detection methods readily determined from the known art,
20 examples of which include, without limitation, immunological techniques (*e.g.*, immunohistochemical staining), fluorescence imaging techniques, and microscopic techniques.

[0066] The inhibitor and factors of the present invention may be contacted with neural stem cells or neural progenitor cells, either *in vitro* or *in vivo* in a subject,
25 by known techniques used for the introduction and administration of proteins, nucleic acids, and other drugs. Examples of methods for contacting the cells with (*i.e.*, treating the cells with) an ATF5 inhibitor or a neurotrophic factor (in protein or nucleic acid form) include, without limitation, absorption, electroporation, immersion, injection, introduction, liposome delivery, transfection, transfusion, vectors, and other protein-delivery
30 and nucleic-acid-delivery vehicles and methods. When target cells are localized to a particular portion of a subject, it may be desirable to introduce the inhibitor and

factors directly to the cells, by injection or by some other means (e.g., by introducing the inhibitor and factors into the blood or another body fluid).

[0067] Where the inhibitor or neurotrophic factor is a protein or other molecule, it may be introduced into a neural stem cell or neural progenitor cell directly, in accordance with conventional techniques and methods disclosed herein. Additionally, a protein inhibitor or factor may be introduced into a neural stem cell or neural progenitor cell indirectly, by introducing into the cell a nucleic acid encoding the inhibitor or factor, in a manner permitting expression of the protein inhibitor or factor. The inhibitor or factor may be introduced into neural stem cells or neural progenitor cells, *in vitro* or *in vivo*, using conventional procedures known in the art, including, without limitation, electroporation, DEAE Dextran transfection, calcium phosphate transfection, monocationic liposome fusion, polycationic liposome fusion, protoplast fusion, creation of an *in vivo* electrical field, DNA-coated microprojectile bombardment, injection with recombinant replication-defective viruses, homologous recombination, *in vivo* gene therapy, *ex vivo* gene therapy, viral vectors, and naked DNA transfer, or any combination thereof. Recombinant viral vectors suitable for gene therapy include, but are not limited to, vectors derived from the genomes of such viruses as retrovirus, HSV, adenovirus, adeno-associated virus, Semiliki Forest virus, cytomegalovirus, lentivirus, and vaccinia virus. The amount of nucleic acid to be used is an amount sufficient to express an amount of protein effective to produce a differentiated neural cell. These amounts may be readily determined by the skilled artisan. It is also within the confines of the present invention that a nucleic acid encoding a protein inhibitor or factor may be introduced into suitable neural stem cells or neural progenitor cells *in vitro*, using conventional procedures, to achieve expression of the protein inhibitor or factor in the cells. Cells expressing protein inhibitor or factor then may be introduced into a subject to produce a differentiated neural cell *in vivo*.

[0068] In accordance with the method of the present invention, ATF5 inhibitors and neurotrophic factors may be administered to a human or animal subject by known procedures, including, without limitation, oral administration, parenteral administration, and transdermal administration. Preferably, the inhibitors or factors are administered parenterally, by intracranial, intraspinal, intrathecal, or subcutaneous injection. The inhibitors and factors of the present invention also may be administered to a subject in accordance with any of the

above-described methods for effecting *in vivo* contact between neural stem cells / neural progenitor cells and ATF5 inhibitors / neurotrophic factors.

[0069] For oral administration, an inhibitor or factor formulation may be presented as capsules, tablets, powders, granules, or as a suspension. The formulation may have
5 conventional additives, such as lactose, mannitol, corn starch, or potato starch. The formulation also may be presented with binders, such as crystalline cellulose, cellulose derivatives, acacia, corn starch, or gelatins. Additionally, the formulation may be presented with disintegrators, such as corn starch, potato starch, or sodium carboxymethylcellulose. The formulation also may be presented with dibasic calcium phosphate anhydrous or sodium
10 starch glycolate. Finally, the formulation may be presented with lubricants, such as talc or magnesium stearate.

[0070] For parenteral administration (*i.e.*, administration by injection through a route other than the alimentary canal), an inhibitor or factor may be combined with a sterile aqueous solution that is preferably isotonic with the blood of the subject. Such a formulation
15 may be prepared by dissolving a solid active ingredient in water containing physiologically-compatible substances, such as sodium chloride, glycine, and the like, and having a buffered pH compatible with physiological conditions, so as to produce an aqueous solution, then rendering said solution sterile. The formulation may be presented in unit or multi-dose containers, such as sealed ampoules or vials. The formulation may be delivered by any mode
20 of injection, including, without limitation, epifascial, intracapsular, intracranial, intracutaneous, intrathecal, intramuscular, intraorbital, intraperitoneal, intraspinal, intrasternal, intravascular, intravenous, parenchymatous, subcutaneous, or sublingual.

[0071] For transdermal administration, an inhibitor or factor may be combined with skin penetration enhancers, such as propylene glycol, polyethylene glycol, isopropanol,
25 ethanol, oleic acid, *N*-methylpyrrolidone, and the like, which increase the permeability of the skin to the inhibitor or factor, and permit the inhibitor or factor to penetrate through the skin and into the bloodstream. The inhibitor/enhancer or factor/enhancer compositions also may be further combined with a polymeric substance, such as ethylcellulose, hydroxypropyl cellulose, ethylene/vinylacetate, polyvinyl pyrrolidone, and the like, to provide the
30 composition in gel form, which may be dissolved in solvent, such as methylene chloride, evaporated to the desired viscosity, and then applied to backing material to provide a patch.

[0072] The present invention provides a method for promoting differentiation of neural stem cells or neural progenitor cells into differentiated neural cells, and for purifying and isolating the neural cells so generated using enhanced green fluorescent protein (eGFP) as a genetic marker. The method described herein for promoting differentiation of neural stem cells or neural progenitor cells *in vitro* provides a source of neurons, or other neural cells of the CNS or PNS, that are available for transplant into a subject. Thus, this method is particularly useful for producing neural cells for use in treating conditions associated with nervous tissue degeneration.

[0073] The term "nervous tissue", as used herein, refers to tissue of the nervous system, which includes the differentiated neural cells of the present invention and progenitors thereof. As further used herein, "nervous tissue degeneration" means a condition of deterioration of nervous tissue, wherein the nervous tissue changes to a lower or less functionally-active form. It is believed that, by promoting differentiation of neural stem cells or neural progenitor cells, the method described herein will be useful in repopulating various injured and/or degenerated nervous tissues in a subject, through production of differentiated neural cells and subsequent transplant thereof into a subject in need of such transplantation.

[0074] Accordingly, the present invention provides a method for treating nervous tissue degeneration in a subject in need of treatment for nervous tissue degeneration, comprising promoting differentiation of neural stem cells or neural progenitor cells into differentiated neural cells, in accordance with the methods described herein, and transplanting the differentiated neural cells into the subject, thereby treating the nervous tissue degeneration. By way of example, the method of the present invention may comprise the following steps: (a) obtaining or generating a culture of neural stem cells or neural progenitor cells; (b) contacting the culture of neural stem cells or neural progenitor cells with an amount of an ATF5 inhibitor effective to produce differentiated neural cells; (c) optionally, contacting the differentiated neural cells with at least one neurotrophic factor; and (d) transplanting the differentiated neural cells into the subject, in an amount effective to treat the nervous tissue degeneration. In one embodiment of the invention, the subject is an embryo. In another embodiment of the invention, the subject is a human. Preferably, the subject has nervous tissue degeneration.

[0075] Nervous tissue degeneration may arise in the CNS or PNS, and may be caused by, or associated with, a variety of disorders, conditions, and factors, including, without limitation, primary neurologic conditions (*e.g.*, neurodegenerative diseases), demyelinating conditions, CNS and PNS traumas and injuries, and acquired secondary effects of non-neural dysfunction (*e.g.*, neural loss secondary to degenerative, pathologic, or traumatic events).
5 Examples of CNS traumas include, without limitation, blunt trauma, hypoxia, and invasive trauma. Examples of acquired secondary effects of non-neural dysfunction include, without limitation, cerebral palsy, congenital hydrocephalus, muscular dystrophy, stroke, and vascular dementia, as well as neural degeneration
10 resulting from any of the following: an injury associated with cerebral hemorrhage, developmental disorders (*e.g.*, a defect of the brain, such as congenital hydrocephalus, or a defect of the spinal cord, such as spina bifida), diabetic encephalopathy, hypertensive encephalopathy, intracranial aneurysms, ischemia, kidney dysfunction, subarachnoid hemorrhage, trauma to the brain
15 and spinal cord, treatment by such therapeutic agents as chemotherapy agents and antiviral agents, vascular lesions of the brain and spinal cord, and other diseases or conditions prone to result in nervous tissue degeneration.

[0076] In one embodiment of the present invention, the nervous tissue degeneration is a peripheral neuropathy in the PNS. As defined herein, the term "peripheral neuropathy"
20 refers to a syndrome of sensory loss, muscle weakness, muscle atrophy, decreased deep-tendon reflexes, and/or vasomotor symptoms. In a subject who has a peripheral neuropathy, myelin sheaths (or Schwann cells) may be primarily affected, or axons may be primarily affected. The peripheral neuropathy may affect a single nerve (mononeuropathy), two or more nerves in separate areas (multiple mononeuropathy), or many nerves simultaneously
25 (polyneuropathy).

[0077] Examples of peripheral neuropathies that may be treated by the methods disclosed herein include, without limitation, peripheral neuropathies associated with acute or chronic inflammatory polyneuropathy, amyotrophic lateral sclerosis (ALS), collagen vascular disorder (*e.g.*, polyarteritis nodosa, rheumatoid arthritis, Sjögren's syndrome, or systemic
30 lupus erythematosus), diphtheria, Guillain-Barré syndrome, hereditary peripheral neuropathy (*e.g.*, Charcot-Marie-Tooth disease (including type I, type II, and all subtypes), hereditary

motor and sensory neuropathy (types I, II, and III, and peroneal muscular atrophy), hereditary neuropathy with liability to pressure palsy (HNPP), infectious disease (*e.g.*, acquired immune deficiency syndrome (AIDS)), Lyme disease (*e.g.*, infection with *Borrelia burgdorferi*), invasion of a microorganism (*e.g.*, leprosy – the leading cause of peripheral neuropathy worldwide, after neural trauma), leukodystrophy, metabolic disease or disorder (*e.g.*, amyloidosis, diabetes mellitus, hypothyroidism, porphyria, sarcoidosis, or uremia), neurofibromatosis, nutritional deficiencies, paraneoplastic disease, peroneal nerve palsy, polio, porphyria, postpolio syndrome, Proteus syndrome, pressure paralysis (*e.g.*, carpal tunnel syndrome), progressive bulbar palsy, radial nerve palsy, spinal muscular atrophy (SMA), a toxic agent (*e.g.*, barbitol, carbon monoxide, chlorobutanol, dapsone, emetine, heavy metals, hexobarbital, lead, nitrofurantoin, orthodinitrophenol, phenytoin, pyridoxine, sulfonamides, triorthocresyl phosphate, the vinca alkaloids, many solvents, other industrial poisons, and certain AIDS drugs (including didanosine and zalcitabine), trauma (including neural trauma – the leading cause of peripheral neuropathy, worldwide), and ulnar nerve palsy (Beers and Berkow, eds., *The Merck Manual of Diagnosis and Therapy*, 17th ed. (Whitehouse Station, NJ: Merck Research Laboratories, 1999) chap. 183). In a preferred embodiment of the present invention, the peripheral neuropathy is ALS or SMA.

[0078] In another embodiment of the present invention, the nervous tissue degeneration is a neurodegenerative disease. Examples of neurodegenerative diseases that may be treated by the methods disclosed herein include, without limitation, Alzheimer's disease, amyotrophic lateral sclerosis (Lou Gehrig's Disease), Binswanger's disease, Huntington's chorea, multiple sclerosis, myasthenia gravis, Parkinson's disease, and Pick's disease.

[0079] It is also within the confines of the present invention for the method described herein to be used to treat nervous tissue degeneration that is associated with a demyelinating condition. Demyelinating conditions are manifested in loss of myelin – the multiple dense layers of lipids and protein which cover many nerve fibers. These layers are provided by oligodendroglia in the CNS, and Schwann cells in the PNS. In patients with demyelinating conditions, demyelination may be irreversible; it is usually accompanied or followed by axonal degeneration, and often by cellular degeneration. Demyelination can occur as a result of neuronal damage or damage to the myelin itself –

whether due to aberrant immune responses, local injury, ischemia, metabolic disorders, toxic agents, or viral infections.

[0080] Central demyelination (demyelination of the CNS) occurs in several conditions, often of uncertain etiology, that have come to be known as the primary demyelinating diseases. Of these, multiple sclerosis is the most prevalent. Other primary demyelinating diseases include adrenoleukodystrophy (ALD), adrenomyeloneuropathy, AIDS-vacuolar myelopathy, HTLV-associated myelopathy, Leber's hereditary optic atrophy, progressive multifocal leukoencephalopathy (PML), subacute sclerosing panencephalitis, and tropical spastic paraparesis. In addition, there are acute conditions in which demyelination can occur in the CNS, *e.g.*, acute disseminated encephalomyelitis (ADEM) and acute viral encephalitis. Furthermore, acute transverse myelitis, a syndrome in which an acute spinal cord transection of unknown cause affects both gray and white matter in one or more adjacent thoracic segments, can also result in demyelination. Finally, there are animal models which mimic features of human demyelinating diseases. Examples include experimental autoimmune neuritis (EAN), demyelination induced by Theiler's virus, and experimental autoimmune encephalomyelitis (EAE) – an autoimmune disease which is experimentally induced in a variety of species and which resembles MS in its clinical and neuropathological aspects.

[0081] The differentiated neural cells of the present invention may be transplanted into a subject in need of treatment by standard procedures known in the art, as well as the methods described herein. By way of example, neural stem cells or neural progenitor cells may be induced with an ATF5 inhibitor, to produce differentiated neural cells. At an appropriate time post-induction (*e.g.*, 3-4 days after induction), the cells may be prepared for transplantation (*e.g.*, partially triturated), and then transplanted into a subject (*e.g.*, into the spinal cord of a chick, HH stage 15-17). To accommodate transplanted tissue, the subject may be suction-lesioned prior to implantation. In one embodiment of the present invention, the differentiated neural cells are transplanted into the spinal cord of a subject, thereby repopulating the subject's spinal cord, and the nervous tissue degeneration is a peripheral neuropathy associated with ALS or SMA.

[0082] In another embodiment of the present invention, the neural stem cells or neural progenitor cells contain an ATF5 transgene that has been engineered to express ATF5 on an

inducible promoter. In this embodiment of the present invention, ATF5 may be expressed in the presence of a suitable inducing agent, thereby permitting propagation of the neural stem cells or neural progenitor cells *in vitro*. Once the cells are transplanted into the subject, however, the inducing agent would be withdrawn, resulting in decreased ATF5 expression, and thereby promoting differentiation of the transplanted cells. Expression of ATF5 would be sustained in the presence of the inducer, and would be shut down once the supply of inducer was depleted (*e.g.*, upon transplant into a subject).

[0083] In an alternative embodiment, a dominant-negative form of ATF5 (an ATF5 inhibitor) may be introduced into the neural stem cells or neural progenitor cells on an inducible promoter. The transgene could be maintained in an uninduced state *in vitro*, permitting propagation of the cells, and then induced with a suitable inducing agent, *in vivo* in a subject, thereby promoting differentiation of the neural stem cells or neural progenitor cells.

[0084] In the method of the present invention, the differentiated neural cells are transplanted into a subject in need of treatment in an amount effective to treat the nervous tissue degeneration. As used herein, the phrase "effective to treat the nervous tissue degeneration" means effective to ameliorate or minimize the clinical impairment or symptoms of the nervous tissue degeneration. For example, where the nervous tissue degeneration is a peripheral neuropathy, the clinical impairment or symptoms of the peripheral neuropathy may be ameliorated or minimized by alleviating vasomotor symptoms, increasing deep-tendon reflexes, reducing muscle atrophy, restoring sensory function, and strengthening muscles. The amount of differentiated neural cells effective to treat nervous tissue degeneration in a subject in need of treatment will vary depending upon the particular factors of each case, including the type of nervous tissue degeneration, the stage of the nervous tissue degeneration, the subject's weight, the severity of the subject's condition, the type of differentiated neural cells, and the method of transplantation. This amount may be readily determined by the skilled artisan, based upon known procedures, including clinical trials, and methods disclosed herein.

[0085] In view of the above-described method for promoting differentiation of neural stem cells and neural progenitor cells into differentiated neural cells, the present invention further provides a method for producing differentiated neural cells, comprising the steps of:

(a) obtaining or generating a culture of neural stem cells or neural progenitor cells; (b) contacting the culture of neural stem cells or neural progenitor cells with an amount of an ATF5 inhibitor effective to produce a subclass of differentiated neural cells; and (c) optionally, contacting the differentiated neural cells with at least one neurotrophic factor.

5 The present invention also provides a population of cells, comprising the differentiated neural cells produced by this method. In one embodiment, some or all of the cells express eGFP.

[0086] In the method of the present invention, any of steps (b)-(c) may be performed *in vitro*, or *in vivo* in a subject. Following any *in vitro* steps, cells may be transplanted into a subject such that the remaining steps are performed *in vivo*. Accordingly, the method of the present invention further comprises the step of transplanting the neural progenitor cells or the differentiated neural cells into a subject. For example, the neural stem cells or neural progenitor cells may contain an ATF5 transgene that has been engineered to express ATF5 on an inducible promoter. In this embodiment of the present invention, ATF5 would be expressed in the presence of a suitable inducing agent, thereby permitting propagation of the neural stem cells or neural progenitor cells *in vitro*. Thereafter, the cells may be transplanted into a subject, such that steps (b) and (c) are carried out *in vivo*. Because the inducing agent would be withdrawn upon transplantation of the cells into the subject, ATF5 expression would be decreased, thereby promoting differentiation of the transplanted cells. Similarly, a culture of neural stem cells or neural progenitor cells may be contacted with an ATF5 inhibitor *in vitro*, to produce differentiated neural cells. The neural cells so produced then may be transplanted into a subject, such that step (c) is carried out *in vivo*. In an alternative method, a culture of neural stem cells or neural progenitor cells may be contacted with an ATF5 inhibitor *in vitro*, to produce differentiated neural cells; and, optionally, the differentiated neural cells may be contacted with at least one neurotrophic factor *in vitro*. The differentiated neural cells then may be transplanted into a subject. In one embodiment of the present invention, the neurons are transplanted into the spinal cord of the subject.

[0087] Because the selective degeneration of specific classes of CNS neurons underlies many neurological disorders, research into the growth, survival, and activity of neurons remains a priority. Unfortunately, however, live neurons are not readily available for such studies. For this reason, the present invention will be of particular importance to researchers in the fields of neuroscience and neurology, as it provides a potentially-unlimited

source of neural cells to be studied. Accordingly, the present invention also provides for uses of the above-described neural progenitor cells and differentiated neural cells in particular areas of research.

[0088] The neural progenitor cells and differentiated neural cells of the present invention will be useful in the analysis of neuron development, function, and death – research which is critical to a complete understanding of neurological diseases. Furthermore, the neural progenitor cells and differentiated neural cells of the present invention will be useful in monitoring synaptic differentiation at sites of contact with target muscles. Finally, the neural progenitor cells and differentiated neural cells of the present invention will facilitate a direct comparison of normal, healthy neurons with degenerated neurons. For such a comparison, both the healthy and the diseased neural cells may be produced using well-known techniques and methods described herein.

[0089] The present invention further provides a method for isolating a pure population of differentiated neural cells and/or purifying a population of differentiated neural cells, comprising the steps of: (a) obtaining or generating a culture of neural stem cells or neural progenitor cells that express enhanced green fluorescent protein (eGFP); (b) contacting the culture of neural stem cells or neural progenitor cells with an amount of an ATF5 inhibitor effective to produce differentiated neural cells, wherein some or all of the differentiated neural cells also express eGFP; (c) optionally, contacting the differentiated neural cells with at least one neurotrophic factor; (d) detecting expression of eGFP in the differentiated neural cells; and (e) isolating the differentiated neural cells that express eGFP. Neural stem cells or neural progenitor cells that express eGFP may be made in accordance with methods disclosed herein.

[0090] According to the method of the present invention, expression of eGFP may be detected in differentiated neural cells by either *in vitro* or *in vivo* assay. As used herein, "expression" refers to the transcription of the eGFP gene into at least one mRNA transcript, or the translation of at least one mRNA into an eGFP protein. The differentiated neural cells may be assayed for eGFP expression by assaying for eGFP protein, eGFP cDNA, or eGFP mRNA. The appropriate form of eGFP will be apparent based on the particular techniques discussed herein.

[0091] Differentiated neural cells may be assayed for eGFP expression, and eGFP expression may be detected in differentiated neural cells, using assays and detection methods well known in the art. Because eGFP provides a non-invasive marker for labeling cells in culture and *in vivo*, expression of eGFP is preferably detected in differentiated neural cells using imaging techniques, particularly bright-field, phase, and fluorescence imaging techniques, as disclosed herein. Differentiated neural cells expressing high levels of eGFP then may be isolated from a cell suspension by sorting (*e.g.*, by FACS sorting, using a Beckman-Coulter Altra flow cytometer), based upon their eGFP fluorescence and forward light scatter, as described below.

[0092] Other methods also may be used to detect eGFP expression in the differentiated neural cells of the present invention. Examples of such detection methods include, without limitation, hybridization analysis, imaging techniques, immunological techniques, immunoprecipitation, radiation detection, Western-blot analysis, and any additional assays or detection methods disclosed herein. For example, differentiated neural cells may be assayed for eGFP expression using an agent reactive with eGFP protein or eGFP nucleic acid. As used herein, "reactive" means the agent has affinity for, binds to, or is directed against eGFP. As further used herein, an "agent" shall include a protein, polypeptide, peptide, nucleic acid (including DNA or RNA), antibody, Fab fragment, F(ab')₂ fragment, molecule, compound, antibiotic, drug, and any combinations thereof. In one embodiment of the present invention, the agent reactive with eGFP is an antibody (*e.g.*, αGFP (Molecular Probes, Inc., Eugene, OR)).

[0093] Following detection of eGFP expression in differentiated neural cells, the extent of eGFP expression in the cells may be measured or quantified, if desired, using one of various quantification assays. Such assays are well known to one of skill in the art, and may include immunohistochemistry, immunocytochemistry, flow cytometry, mass spectroscopy, Western-blot analysis, or an ELISA for measuring amounts of eGFP protein.

[0094] The present invention further provides a method for identifying an agent for use in treating a condition associated with nervous tissue degeneration, as defined above. Examples of conditions associated with nervous tissue degeneration include peripheral neuropathies, demyelinating conditions, and the primary neurologic conditions (*e.g.*, neurodegenerative diseases), CNS and PNS traumas and injuries, and acquired secondary

effects of non-neural dysfunction (*e.g.*, neural loss secondary to degenerative, pathologic, or traumatic events) described herein.

[0095] The method of the present invention comprises the steps of: (a) obtaining or generating a culture of neural stem cells or neural progenitor cells; (b) contacting the culture of cells with an amount of an ATF5 inhibitor effective to produce neurons, wherein some or all of the neurons are degenerated; (c) contacting the degenerated neurons with a candidate agent; and (d) determining if the agent enhances regeneration or survival of some or all of the degenerated neurons. As used herein, the term "enhance regeneration" means augment, improve, or increase partial or full growth (or regrowth) of a neuron (including neurites and the myelin sheath) that has degenerated. As further used herein, the term "growth" refers to an increase in diameter, length, mass, and/or thickness of a neuron (including neurites and the myelin sheath). Regeneration of the neuron may take place in neurons of both the central nervous system and the peripheral nervous system. Additionally, as used herein, the term "enhance survival" of a neuron means increasing the duration of the neuron's viable lifespan, either *in vitro* or *in vivo*. In one embodiment of the present invention, the agent enhances regeneration or survival of degenerated motor neurons.

[0096] In the method of the present invention, degenerated neurons may be contacted with a candidate agent by any of the methods of effecting contact between inhibitors or factors or agents and cells, and any modes of introduction and administration, described herein. Regeneration, and enhanced regeneration, of neurons may be measured or detected by known procedures, including Western blotting for myelin-specific and axon-specific proteins, electron microscopy in conjunction with morphometry, and any of the methods, molecular procedures, and assays known to one of skill in the art. In addition, growth of myelin may be assayed using the *g*-ratio – one measure of the integrity of the axon:myelin association. The *g*-ratio is defined as the axonal diameter divided by the total diameter of the axon and myelin. This ratio provides a reliable measure of relative myelination for an axon of any given size (Bieri *et al.*, Abnormal nerve conduction studies in mice expressing a mutant form of the POU transcription factor, SCIP. *J. Neurosci. Res.*, 50:821-28, 1997). Numerous studies have documented that a *g*-ratio of 0.6

is normal for most fibers (Waxman and Bennett, Relative conduction velocities of small myelinated and nonmyelinated fibres in the central nervous system.

Nature New Biol., 238:217, 1972). In one embodiment of the present invention, the degenerated neurons express enhanced green fluorescent protein (eGFP). It is expected that such neurons will allow for inhibited high-throughput drug screening.

[0097] The inventors have demonstrated herein that neuronal differentiation may be induced by CRE-mediated gene activation, and that such activation is repressed in neural progenitor cells by factors such as ATF5. Accordingly, the present invention further provides a method for suppressing differentiation of neural stem cells or neural progenitor cells into differentiated neural cells, where such cells might otherwise be determined to differentiate. The method of the present invention comprises contacting the neural stem cells or neural progenitor cells with an amount of ATF5, or a peptidomimetic thereof, effective to suppress differentiation in the neural stem cells or neural progenitor cells. This method will permit a pool of these undifferentiated cells to be generated under conditions in which they might otherwise differentiate and cease proliferation. The ATF5 or mimetic may be in the form of a protein, or a nucleic acid encoding the protein, and may be contacted with the cells in accordance with methods previously described.

[0098] The present invention also provides a therapeutic composition, comprising a nucleic acid encoding an ATF5 inhibitor, a vector, and, optionally, a pharmaceutically-acceptable carrier. The pharmaceutically-acceptable carrier must be "acceptable" in the sense of being compatible with the other ingredients of the composition, and not deleterious to the recipient thereof. The pharmaceutically-acceptable carrier employed herein is selected from various organic or inorganic materials that are used as materials for pharmaceutical formulations, and which may be incorporated as analgesic agents, buffers, binders, disintegrants, diluents, emulsifiers, excipients, extenders, glidants, solubilizers, stabilizers, suspending agents, tonicity agents, vehicles, and viscosity-increasing agents. If necessary, pharmaceutical additives, such as antioxidants, aromatics, colorants, flavor-improving agents, preservatives, and sweeteners, may also be added. Examples of acceptable pharmaceutical carriers include carboxymethyl cellulose, crystalline cellulose, glycerin, gum arabic, lactose, magnesium stearate, methyl cellulose, powders, saline, sodium alginate, sucrose, starch, talc, and water, among others.

[0099] The formulations of the present invention may be prepared by methods well-known in the pharmaceutical arts. For example, the ATF5 inhibitor protein or nucleic acid may be brought into association with a carrier or diluent, as a suspension or solution. Optionally, one or more accessory ingredients (*e.g.*, buffers, flavoring agents, surface active agents, and the like) also may be added. The choice of carrier will depend upon the route of administration. The pharmaceutical composition would be useful for administering the ATF5 inhibitor of the present invention to a subject to treat a neural tumor, as discussed below. The ATF5 inhibitor is provided in an amount that is effective to treat the neural tumor in a subject to whom the pharmaceutical composition is administered. That amount may be readily determined by the skilled artisan, as described above.

[00100] As disclosed herein, the inventors have determined that ATF5 expression is elevated in neural tumors, such as neuroblastomas. Therefore, the pharmaceutical composition of the present invention may be useful for treating a neural tumor in a subject. As used herein, the term "tumor" refers to a pathologic proliferation of cells, and includes a neoplasia. The term "neoplasia", as further used herein, refers to the uncontrolled and progressive multiplication of tumor cells under conditions that would not elicit, or would cause cessation of, multiplication of normal cells. Neoplasia results in the formation of a "neoplasm", which is defined herein to mean any new and abnormal growth, particularly a new growth of tissue, in which the growth of cells is uncontrolled and progressive. As used herein, neoplasms include, without limitation, morphological irregularities in cells in tissue of a subject, as well as pathologic proliferation of cells in tissue of a subject, as compared with normal proliferation in the same type of tissue. Additionally, neoplasms include benign tumors and malignant tumors. Malignant neoplasms are distinguished from benign in that the former show a greater degree of anaplasia, or loss of differentiation and orientation of cells, and have the properties of invasion and metastasis. Thus, neoplasia includes "cancer", which herein refers to a proliferation of tumor cells having the unique trait of loss of normal controls, resulting in unregulated growth, lack of differentiation, local tissue invasion, and metastasis.

[00101] Additionally, as used herein, the term "neural tumor" refers to a tumorigenic form of neural cells (*i.e.*, transformed neural cells), and includes astrocytoma cells (*i.e.*, cells of all astrocytomas, including, without limitation, Grades I-IV astrocytomas, anaplastic astrocytoma, astroblastoma, astrocytoma fibrillare, astrocytoma protoplasmaticum, gemistocytic astrocytoma, and glioblastoma multiforme), gliomas, medulloblastomas, neuroblastomas, and other brain tumors. Brain tumors invade and destroy normal tissue, producing such effects as impaired sensorimotor and cognitive function, increased intracranial pressure, cerebral edema, and compression of brain tissue, cranial nerves, and cerebral vessels. Metastases may involve the skull or any intracranial structure. The size, location, rate of growth, and histologic grade of malignancy determine the seriousness of brain tumors. Nonmalignant tumors grow slowly, with few mitoses, no necrosis, and no vascular proliferation. Malignant tumors grow more rapidly, and invade other tissues. However, they rarely spread beyond the CNS, because they cause death by local growth. Drowsiness, lethargy, obtuseness, personality changes, disordered conduct, and impaired mental faculties are the initial symptoms in 25% of patients with malignant brain tumors.

[00102] Brain tumors may be classified by site (*e.g.*, brain stem, cerebellum, cerebrum, cranial nerves, ependyma, meninges, neuroglia, pineal region, pituitary gland, and skull) or by histologic type (*e.g.*, meningioma, primary CNS lymphoma, or astrocytoma). Common primary childhood tumors are cerebellar astrocytomas and medulloblastomas, ependymomas, gliomas of the brain stem, neuroblastomas, and congenital tumors. In adults, primary tumors include meningiomas, schwannomas, and gliomas of the cerebral hemispheres (particularly the malignant glioblastoma multiforme and anaplastic astrocytoma, and the more benign astrocytoma and oligodendroglioma). Overall incidence of intracranial neoplasms is essentially equal in males and females, but cerebellar medulloblastoma and glioblastoma multiforme are more common in males.

[00103] Gliomas are tumors composed of tissue representing neuroglia in any one of its stages of development. They account for 45% of intracranial tumors. Gliomas can encompass all of the primary intrinsic neoplasms of the brain and spinal cord, including astrocytomas, ependymomas, and neurocytomas. Astrocytomas are tumors composed of transformed astrocytes, or astrocytic tumor cells. Such tumors have been classified in order

of increasing malignancy: Grade I consists of fibrillary or protoplasmic astrocytes; Grade II is an astroblastoma, consisting of cells with abundant cytoplasm and two or three nuclei; and Grades III and IV are forms of glioblastoma multiforme, a rapidly growing tumor that is usually confined to the cerebral hemispheres and composed of a mixture of astrocytes, spongioblasts, astroblasts, and other astrocytic tumor cells. Astrocytoma, a primary CNS tumor, is frequently found in the brain stem, cerebellum, and cerebrum. Anaplastic astrocytoma and glioblastoma multiforme are commonly located in the cerebrum.

[00104] In view of the foregoing, the present invention further provides a method for treating a neural tumor in a subject in need of treatment, comprising administering to the subject a pharmaceutical composition comprising an ATF5 inhibitor and a pharmaceutically-acceptable carrier. The ATF5 inhibitor is provided in an amount that is effective to treat the neural tumor in a subject to whom the composition is administered. As used herein, the phrase "effective to treat the neural tumor" means effective to ameliorate or minimize the clinical impairment or symptoms of the neural tumor. For example, the clinical impairment or symptoms of the tumor may be ameliorated or minimized by diminishing any pain or discomfort suffered by the subject; by extending the survival of the subject beyond that which would otherwise be expected in the absence of such treatment; by inhibiting or preventing the development or spread of the tumor; or by limiting, suspending, terminating, or otherwise controlling the maturation and proliferation of cells in the tumor. The amount of ATF5 inhibitor effective to treat a neural tumor in a subject in need of treatment therefor will vary depending upon the particular factors of each case, including the type of neural tumor, the stage of the tumor, the subject's weight, the severity of the subject's condition, and the method of administration. This amount can be readily determined by the skilled artisan. In one embodiment of the present invention, the pharmaceutical composition comprises a nucleic acid encoding an ATF5 inhibitor, a viral vector, and, optionally, a pharmaceutically-acceptable carrier.

[00105] As disclosed herein, according to one proposed pathway, ATF5 binding to CRE DNA-binding sites suppresses differentiation of neural stem cells and neural progenitor cells into differentiated neural cells. Thus, effective ATF5 inhibitors can be designed to replace CRE in its interaction with ATF5. A candidate agent having the ability to bind ATF5

may, as a consequence of this binding, prevent ATF5 binding to CRE through steric hindrance.

[00106] Accordingly, the present invention also provides a method for identifying an agent that inhibits ATF5, by assessing the ability of a candidate agent to inhibit interaction
5 between ATF5 and CRE. The method of the present invention comprises the steps of: (a) contacting a candidate agent with ATF5, in the presence of CRE; and (b) assessing the ability of the candidate agent to inhibit interaction between ATF5 and CRE. An agent that inhibits interaction between ATF5 and CRE may be either natural or synthetic, and may be an agent reactive with ATF5 (*i.e.*, has affinity for, binds to, or is directed against ATF5). An agent
10 that is reactive with ATF5, as disclosed herein, may have the ability to inhibit interaction between ATF5 and CRE by binding to ATF5. A candidate agent having the ability to bind to ATF5 may, as a consequence of this binding, inhibit ATF5 activity through steric interactions (without binding to CRE itself). A CRE-luciferase reporter assay may be used to gauge such interactions, as described herein (Example 11).

15 [00107] In accordance with the method of the present invention, a CRE-like agent that binds ATF5 may be identified using an *in vitro* assay (*e.g.*, a direct binding assay, competitive binding assay, etc.). In a direct binding assay, for example, the binding of a candidate agent to ATF5 or a peptide fragment thereof may be measured directly. A candidate agent may be supplied by a peptide library, for example. Alternatively, in a
20 competitive binding assay, standard methodologies may be used in order to assess the ability of a candidate agent to bind ATF5, and thereby inhibit CRE-ATF5 interaction. In such a competitive binding assay, the candidate agent competes with CRE for binding to ATF5 (but does not bind directly to CRE). Once bound to ATF5, the candidate agent could sterically hinder binding of CRE to ATF5, thereby preventing interaction between CRE and ATF5. A
25 competitive binding assay represents a convenient way to assess inhibition of CRE-ATF5 interaction, since it allows the use of crude extracts containing ATF5 and CRE.

[00108] A competitive binding assay may be carried out by adding ATF5, or an extract containing ATF5 biological activity (as defined above), to a mixture containing the candidate agent and labeled CRE, both of which are present in the mixture in known concentrations.
30 After incubation, the ATF5-agent complex may be separated from the unbound labeled CRE and unlabeled candidate agent, and counted. The concentration of the candidate agent

required to inhibit 50% of the binding of the labeled CRE to ATF5 (IC_{50}) then may be calculated.

[00109] The binding assay formats described herein employ labeled assay components. Labeling of CRE or ATF5 may be accomplished using one of a variety of different chemiluminescent and radioactive labels known in the art. The label of the present invention may be, for example, a nonradioactive or fluorescent marker, such as biotin, fluorescein (FITC), acridine, cholesterol, or carboxy-X-rhodamine, which can be detected using fluorescence and other imaging techniques readily known in the art. Alternatively, the label may be a radioactive marker, including, for example, a radioisotope. The radioisotope may be any isotope that emits detectable radiation, including, without limitation, ^{35}S , ^{32}P , ^{125}I , 3H , or ^{14}C . The label may also be luciferase, for use in a CRE-luciferase reporter assay, as described below (Example 11).

[00110] Qualitative results of the above-described assays may be obtained by competitive autoradiographic-plate binding assays; alternatively, Scatchard plots may be used to generate quantitative results. The labels of the present invention may be coupled directly or indirectly to the desired component of the assay, according to methods well known in the art. The choice of label depends on a number of relevant factors, including the sensitivity required, the ease of conjugation with the compound to be labeled, stability requirements, and available instrumentation.

[00111] Both direct and competitive binding assays may be used in a variety of different configurations. In one competitive binding assay, for example, the candidate agent may compete against labeled CRE (the labeled analyte) for a specific binding site on ATF5 (the capture agent) that is bound to a solid substrate, such as a column chromatography matrix or tube. Alternatively, the candidate agent may compete for a specific binding site on labeled ATF5 (the labeled analyte) against wild-type CRE or a fragment thereof (the capture agent) that is bound to a solid substrate. The capture agent is bound to the solid substrate in order to effect separation of bound labeled analyte from the unbound labeled analyte. In either type of competitive binding assay, the concentration of labeled analyte that binds the capture agent bound to the solid substrate is inversely proportional to the ability of a candidate agent to compete in the binding assay. The amount of inhibition of labeled analyte

by the candidate agent depends on the binding assay conditions and on the concentrations of candidate agent, labeled analyte, and capture agent that are used.

[00112] Another competitive binding assay, for use in detecting agents that bind to ATF5, may be conducted in a liquid phase. In this type of assay, any of a variety of techniques known in the art may be used to separate the bound labeled analyte (which may be either CRE or ATF5) from the unbound labeled analyte. Following such separation, the amount of bound labeled analyte may be determined. The amount of unbound labeled analyte present in the separated sample is inversely proportional to the amount of bound labeled analyte.

10 [00113] In the further alternative, a homogeneous binding assay may be performed, in which a separation step is not needed. In this type of binding assay, the label on the labeled analyte (which may be either CRE or ATF5) is altered by the binding of the analyte to the capture agent. This alteration in the labeled analyte results in a decrease or increase in the signal emitted by the label, so that measurement of the label at the end of the binding assay allows for detection or quantification of the analyte.

[00114] Under specified assay conditions, a candidate agent is considered to be capable of inhibiting the binding of CRE to ATF5 in a competitive binding assay if the amount of binding of the labeled analyte to the capture agent is decreased by 50% or more (preferably 90% or more). Where a direct binding assay configuration is used, a candidate agent is considered to bind ATF5 when the signal measured is twice the background level or higher. Furthermore, as proof of the specificity of the candidate agent identified using an ATF5 competitive binding assay, binding competition also may be performed using purified ATF5 in the presence of washed ribosomes. A functional assay, such as a luciferase assay, also may be used to screen for ATF5 inhibitors, as described herein.

25 [00115] As disclosed herein, ATF5 has been implicated in a number of biological events in neural stem cells, neural progenitor cells, and neuroblastoma cells. For example, it has been shown that ATF5 plays a role in the differentiation of neural stem and progenitor cells, and may be associated with uncontrolled cell proliferation in neuroblastomas and other neural tumors. Accordingly, it is clear that therapeutics designed to inhibit ATF5 (*i.e.*, those which bind to, or are otherwise reactive with, ATF5) may be useful in regulation of a number

30

of ATF5-associated biological events, including differentiation of neural stem cells and neural progenitor cells, and control of proliferation of neural tumor cells.

[00116] Thus, once the candidate agent of the present invention has been screened, and has been determined to have a suitable inhibitory effect on ATF5 (i.e., it is reactive with ATF5, it binds ATF5, or it otherwise inactivates ATF5), it may be evaluated for its effect on differentiation of neural stem cells or neural progenitor cells, or on tumor-cell proliferation. In particular, the candidate agent may be assessed for its ability to act as a promoter of differentiation, or as an inhibitor of tumor-cell division proliferation, or to otherwise function as an appropriate tumor-suppressing agent. It is expected that the ATF5 inhibitor of the present invention will be useful for promoting differentiation of neural stem cells and neural progenitor cells, and for treating neural tumors, including those disclosed herein. Furthermore, the inventors propose that the ATF5 inhibitor of the present invention might be useful for restoring proliferation control in tumor cells

[00117] Accordingly, the present invention further comprises the steps of: (c) contacting the candidate agent with neural stem cells or neural progenitor cells containing ATF5; and (d) determining if the agent has an effect on an ATF5-associated biological event in the neural stem cells or neural progenitor cells. As used herein, an "ATF5-associated biological event" includes a biochemical or physiological process in which ATF5 levels or activity have been implicated. As disclosed herein, examples of ATF5-associated biological events include, without limitation, binding to, and interaction with, CRE; regulation of differentiation in neural stem cells or neural progenitor cells; and proliferation of neural tumor cells. As further used herein, a cell "containing ATF5" is a cell in which ATF5, or a derivative or homologue thereof, is naturally expressed or naturally occurs.

[00118] According to this method of the present invention, a candidate agent may be contacted with one or more neural stem cells or neural progenitor cells *in vitro*. For example, a culture of cells may be incubated with a preparation containing the candidate agent. The candidate agent's effect on an ATF5-

associated biological event then may be assessed by any biological assays or methods known in the art, including histological analyses. In one embodiment of the present invention, the neural stem cells or neural progenitor cells express luciferase (see Examples 3 and 11).

5 [00119] The present invention is further directed to agents identified by the above-described identification methods. Such agents may be useful for promoting differentiation of neural stem cells or neural progenitor cells, and for treating an ATF5-associated condition. As used herein, an "ATF5-associated condition" is a condition, disease, or disorder in which ATF5 levels or activity have been implicated, and includes the following: an ATF5-
10 associated biological event, and neural tumors. The ATF5-associated condition may be treated in the subject by administering to the subject an amount of the agent effective to treat the ATF5-associated condition in the subject. This amount may be readily determined by one skilled in the art.

[00120] Accordingly, in one embodiment, the present invention provides a method for
15 promoting differentiation in neural stem cells or neural progenitor cells, by contacting the cells with the above-described agent, in an amount effective to promote differentiation in the cells. In another embodiment, the present invention provides a method for treating or preventing a neural tumor in a subject, by administering to the subject the above-described agent, in an amount effective to treat or prevent the neural tumor in the subject. In a
20 preferred embodiment of the present invention, the neural tumor is a neuroblastoma.

[00121] The present invention also provides a pharmaceutical composition comprising the agent identified by the above-described identification method and a pharmaceutically-acceptable carrier. Examples of suitable pharmaceutically-acceptable carriers, and methods of preparing pharmaceutical formulations and compositions, are described above. The
25 pharmaceutical composition of the present invention would be useful for contacting neural stem cells or neural progenitor cells with an agent that inhibits interaction between CRE and ATF5, in order to promote differentiation of the cells, and would also be useful for treating an ATF5-associated condition. In such a case, the pharmaceutical composition is administered to a subject in an
30 amount effective to treat the ATF5-associated condition.

[00122] The inventors have demonstrated herein that ATF5 expression is elevated in neuroblastoma cells. Thus, ATF5 represents a marker for neuroblastoma. Accordingly, the present invention further provides a method for determining whether a subject has a neural tumor, thereby permitting the diagnosis of such a neural tumor in the subject. The subject
5 may be any of those described above. Preferably, the subject is a human. Examples of neural tumors have been previously discussed. In one embodiment of the present invention, the neural tumor is a neuroblastoma. The method of the present invention comprises assaying a diagnostic sample of the subject for ATF5, wherein detection of an ATF5 level elevated above normal is diagnostic of a neural tumor in the subject. As used herein, "ATF5" includes
10 both an ATF5 protein and an ATF5 analogue, as discussed above.

[00123] In accordance with the method of the present invention, the diagnostic sample of a subject may be assayed *in vitro* or *in vivo*. Where the assay is performed *in vitro*, a diagnostic sample from the subject may be removed using standard procedures. The diagnostic sample may be any nervous tissue, including brain tissue, which may be removed
15 by standard biopsy. In addition, the diagnostic sample may be any tissue known to have a neural tumor, any tissue suspected of having a neural tumor, or any tissue believed not to have a neural tumor. In a preferred embodiment of the present invention, the diagnostic sample contains post-mitotic cells. More preferably, the diagnostic sample contains neural-tumor cells.

[00124] Protein may be isolated and purified from the diagnostic sample of the present invention using standard methods known in the art, including, without limitation, extraction from a tissue (*e.g.*, with a detergent that solubilizes the protein) where necessary, followed by affinity purification on a column, chromatography (*e.g.*, FTLC and HPLC), immuno-precipitation (with an antibody to ATF5), and precipitation (*e.g.*, with isopropanol and a
25 reagent such as Trizol). Isolation and purification of the protein may be followed by electrophoresis (*e.g.*, on an SDS-polyacrylamide gel). Nucleic acid may be isolated from a diagnostic sample using standard techniques known to one of skill in the art.

[00125] In accordance with the method of the present invention, a neural tumor in a subject is diagnosed by assaying a diagnostic sample of the subject for ATF5. The level of
30 ATF5 in the sample, for example, may be detected by measuring ATF5 amounts in the sample. A diagnostic sample may be assayed for the level of ATF5 by assaying for

ATF5 protein, ATF5 cDNA, or ATF5 mRNA. The appropriate form of ATF5 will be apparent based on the particular techniques discussed herein. Preferably, the diagnostic sample of the present invention is assayed for the level of ATF5 protein. It is contemplated that the diagnostic sample may be assayed for
5 expression of any or all forms of ATF5 protein (including precursor forms, endoproteolytically-processed forms, the 22-kDa and 30-kDa forms, and other forms resulting from post-translational modification) in order to determine whether a subject or patient has a neural tumor.

[00126] Alternatively, the level of ATF5 in the sample may be detected by detecting
10 above-normal interaction of ATF5 and CRE. Accordingly, in one embodiment of the present invention, the level of ATF5 elevated above normal is detected by detecting above-normal interaction of ATF5 and CRE. Methods for detecting interaction between CRE and ATF5 have been discussed above.

[00127] As used herein, the term "elevated above normal" means that ATF5 is detected
15 at a level that is significantly greater than the level expected for the same type of diagnostic sample taken from a nondiseased subject or patient (*i.e.*, one who does not have a neural tumor) of the same gender and of similar age. As further used herein, "significantly greater" means that the difference between the level of ATF5 that is elevated above normal, and the expected (normal) level of ATF5, is of statistical significance. Preferably, the level of ATF5
20 elevated above normal is a level that is at least 10% greater than the level of ATF5 otherwise expected in the diagnostic sample. Where ATF5 is expected to be absent from a particular diagnostic sample taken from a particular subject or patient, the normal level of ATF5 for that subject or patient is nil. Where a particular diagnostic sample taken from a particular subject or patient is expected to have a low, constitutive level of ATF5, that low level is the normal
25 level of ATF5 for that subject or patient. As disclosed herein, ATF5 is generally present at lower levels in post-mitotic neurons, than in neural stem cells, neural progenitor cells, or neural tumor cells.

[00128] Expected or normal levels of ATF5 for a particular diagnostic sample taken from a subject or patient may be easily determined by assaying nondiseased subjects of a
30 similar age and of the same gender. For example, diagnostic samples may be obtained from at least 30 normal, healthy men between the ages of 25 and 80, to determine the normal

quantity of ATF5 in males. A similar procedure may be followed to determine the normal quantity of ATF5 in females. Once the necessary or desired samples have been obtained, the normal quantity of ATF5 in men and women may be determined using a standard assay for quantification, such as flow cytometry, Western-blot analysis, or an ELISA for measuring protein quantities, as described below. For example, an ELISA may be run on each sample in duplicate, and the mean and standard deviation of the quantity of ATF5 may be determined. If necessary, additional subjects may be recruited before the normal quantity of ATF5 is determined. A similar type of procedure may be used to determine the expected or normal level of interaction between ATF5 and CRE for a particular diagnostic sample taken from a subject or patient.

[00129] In accordance with the method of the present invention, a diagnostic sample of a subject may be assayed for ATF5 (or for interaction between ATF5 and CRE), and ATF5 (or interaction between ATF5 and CRE) may be detected in a diagnostic sample, using assays and detection methods readily determined from the known art (*e.g.*, immunological techniques, hybridization analysis, fluorescence imaging techniques, and/or radiation detection, etc.), as well as any assays and detection methods disclosed herein (*e.g.*, immunoprecipitation, Western-blot analysis, etc.). For example, a diagnostic sample of a subject may be assayed for ATF5 using an agent reactive with ATF5. The agent may include any of those described above. Preferably, the agent of the present invention is labeled with a detectable marker or label.

[00130] In one embodiment of the present invention, the agent reactive with ATF5 is an antibody. As used herein, the antibody of the present invention may be polyclonal or monoclonal. In addition, the antibody of the present invention may be produced by techniques well known to those skilled in the art. Polyclonal antibody, for example, may be produced by immunizing a mouse, rabbit, or rat with purified ATF5 or with a short peptide sequence thereof. Monoclonal antibody then may be produced by removing the spleen from the immunized mouse, and fusing the spleen cells with myeloma cells to form a hybridoma which, when grown in culture, will produce a monoclonal antibody.

[00131] The antibodies used herein may be labeled with a detectable marker or label. Labeling of an antibody may be accomplished using one of a variety of labeling techniques,

including peroxidase, chemiluminescent labels known in the art, and radioactive labels known in the art. The detectable marker or label of the present invention may be, for example, a nonradioactive or fluorescent marker, such as biotin, fluorescein (FITC), acridine, cholesterol, or carboxy-X-rhodamine, which can be detected using fluorescence and other
5 imaging techniques readily known in the art. Alternatively, the detectable marker or label may be a radioactive marker, including, for example, a radioisotope. The radioisotope may be any isotope that emits detectable radiation, such as ^{35}S , ^{32}P , ^{125}I , ^3H , or ^{14}C . Radioactivity emitted by the radioisotope can be detected by techniques well known in the art. For example, gamma emission from the radioisotope may be detected using gamma imaging
10 techniques, particularly scintigraphic imaging. Preferably, the agent of the present invention is a high-affinity antibody labeled with a detectable marker or label.

[00132] Where the agent of the present invention is an antibody reactive with ATF5, a diagnostic sample taken from the subject may be purified by passage through an affinity column which contains an anti-ATF5 antibody as a ligand attached to a solid support, such as
15 an insoluble organic polymer in the form of a bead, gel, or plate. The antibody attached to the solid support may be used in the form of a column. Examples of suitable solid supports include, without limitation, agarose, cellulose, dextran, polyacrylamide, polystyrene, sepharose, or other insoluble organic polymers. The antibody may be further attached to the solid support through a spacer molecule, if desired. Appropriate binding conditions (*e.g.*,
20 temperature, pH, and salt concentration) for ensuring binding of the agent and the antibody may be readily determined by the skilled artisan. In a preferred embodiment, the antibody is attached to a sepharose column, such as Sepharose 4B.

[00133] Where the agent is an antibody, a diagnostic sample of the subject may be assayed for ATF5 using binding studies that utilize one or more antibodies immunoreactive
25 with ATF5, along with standard immunological detection techniques. For example, the ATF5 protein eluted from the affinity column may be subjected to an ELISA assay, Western-blot analysis, flow cytometry, or any other immunostaining method employing an antigen-antibody interaction. Preferably, the diagnostic sample is assayed for ATF5 using Western blotting.

30 [00134] Alternatively, a diagnostic sample of a subject may be assayed for ATF5 using hybridization analysis of nucleic acid extracted from the diagnostic sample taken from the

subject. According to this method of the present invention, the hybridization analysis may be conducted using Northern blot analysis of mRNA. This method also may be conducted by performing a Southern blot analysis of DNA using one or more nucleic acid probes, which hybridize to nucleic acid encoding ATF5. The nucleic acid probes may be prepared by a variety of techniques known to those skilled in the art, including, without limitation, the following: restriction enzyme digestion of ATF5 nucleic acid; and automated synthesis of oligonucleotides having sequences which correspond to selected portions of the nucleotide sequence of the ATF5 nucleic acid, using commercially-available oligonucleotide synthesizers, such as the Applied Biosystems Model 392 DNA/RNA synthesizer.

[00135] The nucleic acid probes used in the present invention may be DNA or RNA, and may vary in length from about 8 nucleotides to the entire length of the ATF5 nucleic acid. The ATF5 nucleic acid used in the probes may be derived from mammalian ATF5. The nucleotide sequence for human ATF5, for example, is known. Using this sequence as a probe, the skilled artisan could readily clone a corresponding ATF5 cDNA from other species. In addition, the nucleic acid probes of the present invention may be labeled with one or more detectable markers or labels. Labeling of the nucleic acid probes may be accomplished using one of a number of methods known in the art – *e.g.*, nick translation, end labeling, fill-in end labeling, polynucleotide kinase exchange reaction, random priming, or SP6 polymerase (for riboprobe preparation) – along with one of a variety of labels – *e.g.*, radioactive labels, such as ³⁵S, ³²P, or ³H, or nonradioactive labels, such as biotin, fluorescein (FITC), acridine, cholesterol, or carboxy-X-rhodamine (ROX). Combinations of two or more nucleic acid probes (or primers), corresponding to different or overlapping regions of the ATF5 nucleic acid, also may be used to assay a diagnostic sample for ATF5, using, for example, PCR or RT-PCR.

[00136] The detection of ATF5 (or interaction between ATF5 and CRE) in the method of the present invention may be followed by an assay to measure or quantify the extent of ATF5 in a diagnostic sample of a subject. Such assays are well known to one of skill in the art, and may include immunohistochemistry/immunocytochemistry, flow cytometry, mass spectroscopy, Western-blot analysis, or an ELISA for measuring amounts of ATF5 protein. For example, to use an immunohistochemistry assay, histological (paraffin-embedded) sections of tissue may be placed on slides, and then incubated with an antibody against

ATF5. The slides then may be incubated with a second antibody (against the primary antibody), which is tagged to a dye or other colorimetric system (*e.g.*, a fluorochrome, a radioactive agent, or an agent having high electron-scanning capacity), to permit visualization of ATF5 present in the sections.

5 [00137] It is contemplated that the diagnostic sample in the present invention frequently will be assayed for ATF5 (or interaction between ATF5 and CRE) not by the subject or patient, nor by his/her consulting physician, but by a laboratory technician or other clinician. Accordingly, the method of the present invention further comprises providing to a subject's or patient's consulting physician a report of the results obtained upon assaying a
10 diagnostic sample of the subject or patient for ATF5.

[00138] The present invention further provides a method for assessing the efficacy of therapy to treat a neural tumor in a subject or patient who has undergone or is undergoing treatment for a neural tumor. The method of the present invention comprises assaying a diagnostic sample of the subject or patient for ATF5, wherein a normal level of ATF5 in the
15 diagnostic sample is indicative of successful therapy to treat a neural tumor, and a level of ATF5 elevated above normal in the diagnostic sample is indicative of a need to continue therapy to treat a neural tumor. In one embodiment of the present invention, a level of ATF5 elevated above normal is detected by detecting above-normal interaction between ATF5 and CRE. The neural tumor may be any of those described above. The diagnostic sample may be
20 assayed for ATF5 (or interaction between ATF5 and CRE) *in vitro* or *in vivo*. In addition, the diagnostic sample may be assayed for ATF5 (or interaction between ATF5 and CRE) using all of the various assays and methods of detection and quantification described above. This method of the present invention provides a means for monitoring the effectiveness of therapy to treat a neural tumor by permitting the periodic assessment of levels of ATF5 (or interaction
25 between ATF5 and CRE) in a diagnostic sample taken from a subject or patient.

[00139] According to the method of the present invention, a diagnostic sample of a subject or patient may be assayed, and levels of ATF5 (or interaction between ATF5 and CRE) may be assessed, at any time following the initiation of therapy to treat a neural tumor. For example, levels of ATF5 (or interaction between ATF5 and CRE) may be assessed while
30 the subject or patient is still undergoing treatment for a neural tumor. Where levels of ATF5 detected in an assayed diagnostic sample of the subject or patient continue to remain elevated

above normal, a physician may choose to continue with the subject's or patient's treatment for the neural tumor. Where levels of ATF5 in an assayed diagnostic sample of the subject or patient decrease through successive assessments, it may be an indication that the treatment for a neural tumor is working, and that treatment doses could be decreased or even ceased.

5 Where levels of ATF5 in an assayed diagnostic sample of the subject or patient do not rapidly decrease through successive assessments, it may be an indication that the treatment for a neural tumor is not working, and that treatment doses could be increased. Where ATF5 is no longer detected in an assayed diagnostic sample of a subject or patient at a level elevated above normal, a physician may conclude that the treatment for a neural tumor has been
10 successful, and that such treatment may cease.

[00140] It is within the confines of the present invention to assess levels of ATF5 (or interaction between ATF5 and CRE) following completion of a subject's or patient's treatment for a neural tumor, in order to determine whether the neural tumor has recurred in the subject or patient. Accordingly, an assessment of levels of ATF5 (or interaction between
15 ATF5 and CRE) in an assayed diagnostic sample may provide a convenient way to conduct follow-ups of patients who have been diagnosed with a neural tumors. Furthermore, it is within the confines of the present invention to use assessed levels of ATF5 (or interaction between ATF5 and CRE) in an assayed diagnostic sample as a clinical or pathologic staging tool, as a means of determining the extent of a neural tumor in the subject or patient, and as a
20 means of ascertaining appropriate treatment options.

[00141] A correlation exists, in general, between levels of ATF5 in post-mitotic neural cells and neuroblastoma. Therefore, it is also contemplated in the present invention that assaying a diagnostic sample of a subject for ATF5 may be a useful means of providing information concerning the prognosis of a subject or patient who has a neural tumor.

25 Accordingly, the present invention further provides a method for assessing the prognosis of a subject who has a neural tumor, comprising assaying a diagnostic sample of the subject for ATF5, wherein the subject's prognosis improves with a decreased level of ATF5 in the diagnostic sample, and the subject's prognosis worsens with an increased level of ATF5 in the diagnostic sample. In one embodiment of the present invention, the level of ATF5 elevated
30 above normal is detected by detecting above-normal interaction between ATF5 and CRE. Suitable diagnostic samples, assays, and detection and quantification methods for use in the

method of the present invention have already been described. This method of the present invention provides a means for determining the prognosis of a subject or patient diagnosed with a neural tumor based upon the level of ATF5, or interaction between ATF5 and CRE, in an assayed diagnostic sample of the subject or patient.

5. [00142] According to the method of the present invention, a diagnostic sample of a subject or patient may be assayed, and levels of ATF5 (or interaction between ATF5 and CRE) may be assessed, at any time during or following the diagnosis of a neural tumor in the subject or patient. For example, levels of ATF5 (or interaction between ATF5 and CRE) in an assayed diagnostic sample may be assessed before the subject or patient undergoes
10 treatment for a neural tumor, in order to determine the subject's or patient's initial prognosis. Additionally, levels of ATF5 (or interaction between ATF5 and CRE) in an assayed diagnostic sample may be assessed while the subject or patient is undergoing treatment for a neural tumor, in order to determine whether the subject's or patient's prognosis has become more or less favorable through the course of treatment.

15 [00143] For example, where the level of ATF5 detected in an assayed diagnostic sample of the subject or patient is, or continues to remain, significantly high, a physician may conclude that the subject's or patient's prognosis is unfavorable. Where the level of ATF5 in an assayed diagnostic sample of the subject or patient decreases through successive assessments, it may be an indication that the subject's or patient's prognosis is improving.
20 Where the level of ATF5 in an assayed diagnostic sample of the subject or patient does not decrease significantly through successive assessments, it may be an indication that the subject's or patient's prognosis is not improving. Finally, where the level of ATF5 is low, or is normal, in a diagnostic sample of the subject or patient, a physician may conclude that the subject's or patient's prognosis is favorable.

25 [00144] The discovery that ATF5 can be detected in neural tumor cells provides a means of identifying patients with a neural tumor, and presents the potential for commercial application in the form of a test for the diagnosis of a neural tumor. The development of such a test could provide general screening procedures. Such procedures can assist in the early detection and diagnosis of a neural tumor, and can provide a method for the follow-up of
30 patients in whom a level of ATF5 elevated above normal has been detected.

[00145] Accordingly, the present invention further provides a kit for use as an assay of a neural tumor, comprising an ATF5-specific agent and reagents suitable for detecting ATF5. The ATF5-specific agent may be any agent reactive with ATF5 protein or nucleic acid, including a nucleic acid probe which hybridizes to nucleic acid encoding ATF5, an antibody, and any of the agents described above. The agent may be used in any of the above-described assays or methods for detecting or quantifying levels of ATF5. Preferably, the agent of the present invention is labeled with a detectable marker or label.

[00146] The present invention is described in the following Examples, which are set forth to aid in the understanding of the invention, and should not be construed to limit in any way the scope of the invention as defined in the claims which follow thereafter.

EXAMPLES

EXAMPLE 1 – REAGENTS

[00147] Cell-culture media, RPMI 1640 and DMEM, and molecular biology reagents, Taq platinum DNA polymerase, SuperScript II reverse transcriptase, and LipofectAMINE 2000, were obtained from Invitrogen, Inc. (Carlsbad, CA). Donor horse and fetal bovine serum were obtained from JRH Biosciences, Inc. (Lenexa, KA). The Marathon cDNA amplification library kit was obtained from Clontech (Palo Alto, CA), and PCR primers were obtained from Integrated DNA Technologies or Life Technologies, Inc. Anti-FLAG M2 antibody was from Sigma Corp. (St. Louis, MO).

EXAMPLE 2 – CELL CULTURE

[00148] PC12 cells were grown on collagen-coated dishes, as previously described (Greene *et al.*, Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor. *Proc. Natl. Acad. Sci. U S A*, 73:2424-28, 1998), with or without human recombinant nerve growth factor (NGF) (Genentech, Inc.). Dissociated cultures of telencephalic cells were prepared from E14 Sprague-Dawley rats. Telencephalic cells were trypsinized (0.05% in 0.53 mM EDTA; Invitrogen, Inc.) for 30 min (Li *et al.*, Neuronal differentiation of precursors in the neocortical ventricular zone is triggered by BMP. *J. Neurosci.*, 18:8853-62, 1998), and dissociated cells were centrifuged and re-suspended in DMEM containing 5% FBS, 10 ng/ml EGF, and 20 ng/ml bFGF, then plated on 24-well dishes coated with polylysine at $3\text{-}5 \times 10^5$ cells per well (Laywell *et al.*,

Multipotent neurospheres can be derived from forebrain subependymal zone and spinal cord of adult mice after protracted postmortem intervals. *Exp. Neurol.*, 156:430-33, 1999). The presence of bFGF promotes proliferation of the progenitor cells, but does not interfere with their differentiation into neurons (Ghosh and Greenberg, Distinct roles for bFGF and NT-3 in the regulation of cortical neurogenesis. *Neuron*, 15:89-03, 1995).

[00149] Adherent clonal neurosphere cultures were prepared from newborn mouse subependymal zone cells, as previously described (Kukekov *et al.*, A nestin-negative precursor cell from the adult mouse brain gives rise to neurons and glia. *Glia*, 21:399-07, 1997; Kukekov *et al.*, Multipotent stem/progenitor cells with similar properties arise from two neurogenic regions of adult human brain. *Exp. Neurol.*, 156:333-44, 1999). The cell suspension used to generate neurospheres was filtered through sterile gauze, and visually verified to contain only single cells.

EXAMPLE 3 – CLONING OF FULL-LENGTH rATF5 AND PLASMID CONSTRUCTS

[00150] SAGE tag, CATGAGAACCTAGTC, was found in rat EST, UI-R-G0-ur-g-10-0-UI (GenBankTM/EBI accession number AI576016), which, in turn, showed high homology with the 3' end of murine ATF5. To clone the open-reading frame of rat ATF5, PCR antisense primer 5'-CTTGGTTTCTCAGTTGCAC-3' (derived from the sequence of the above EST) was used for 5' RACE PCR, using the Clontech Marathon kit according to the manufacturer's protocol. The first-strand cDNA PCR template was prepared from 5 µg of PC12 cell total RNA, by reverse transcription with Superscript II. The products of the 5' RACE PCR included the second of 2 potential Kozak start sites.

[00151] Cloning of the rATF5 open-reading frame that included the first potential start site was achieved with sense PCR primer, 5'-TGCACCTGTGCCTCAGCCATGTC-3'. This sequence was obtained from an EST sequence (GenBankTM/EBI accession number AW917099) that overlapped with the 5' end of the 5' RACE PCR product described above. Both potential rATF5 forms were FLAG-tagged, by PCR, with sense primers, 5'-CTCGAGA ACCATGGACTACAAGGACGATGATGACAAAGGATCACTCCTGGCGACCCT-3', and 5'-CTCGAGAAGCATGGACTACAAGGACGATGATGACAAAGGAGCATCCCTACTC AAGAA-3'. 5'-GAATTCTCGAGCTTGGTTTCTCAGTTGCAC-3' was the antisense primer for both ATF5s. NTazip-ATF5 was constructed by overlapping PCR, using FLAG-tagged ATF5 (potential start site 2 form) as the template.

[00152] PCR product 1 was produced with 5'-CTCGAGAAGCATGGACTACAA
 GGACGATGATGACAAAGGAGCATCCCTACTCAAGAA-3' and 5'-TTCTTCTGCT
 TCTTTTTCTAGTAGTTCTTCGTTTTCTCTTGCTAGTTCTTCTGCTCTTTGTTGAGG
 GTGCTGGCAGGACTAGGATA-3' as primers, and PCR product 2 was made with 5'-GC
 5 AAGAGAAAACGAAGAACTACTAGAAAAAGAAGCAGAAGAACTAGAACAAAG
 AAATGCAGAGCTAGAGGGCGAGTGCCAAGGG-3' and 5'-GAATTCTCGAGCTTG
 GTTCTCAGTTGCAC-3' as primers. Products 1 and 2 were mixed, and the product (FL-
 NTazip-ATF5) was PCR amplified with 5'-CTCGAGAAGCATGGACTACAAGGACGAT
 GATGACAAAGGAGCATCCCTACTCAAGAA-3' and 5'-GAATTCTCGAGCTTGGTTT
 10 CTCAGTTGCAC-3'. To generate NTazip-ATF5, the activation domain was removed from
 FL-NTazip-ATF5 by PCR, using primers 5'-GAATTCAACCATGGACTACAAGGACGA
 TGATGACAAAATGGCATCTATGACTGGAGGACAACAAATGGGAAGAGACCCA
 GACCTCGAACAAAGAGCAGAA-3' (sense) and 5'-GAATTCTCGAGCTTGGTTTCTCA
 GTTGCAC-3' (antisense).

15 [00153] NTazip-ATF5 was N-terminal FLAG-tagged with a predicted open-reading
 frame of MDYKDDDDKMASMTGGQQMGRDPDLE**QRAEELRENEELLEKEAEELE**
QENAELEGECQGLEARNREL RERAESVEREIQYVKDLLIEVYKARSQRTRSA, where
 the DNA binding motif was replaced with an amphipathic acidic α -helical sequence, as
 marked in bold (Moll *et al.*, Attractive interhelical electrostatic interactions in the proline-
 20 and acidic-rich region (PAR) leucine zipper subfamily preclude heterodimerization with other
 basic leucine zipper subfamilies. *J. Biol. Chem.*, 275:34826-832, 2000). All PCR products
 were subcloned into the Topo II pCR 2.1 vector, and were sequenced to verify identity. After
 confirmation, all full-length constructs were subcloned into the EcoR1 sites of the pCMS-
 eGFP vector.

25 [00154] Retrovirus plasmids were constructed by blunt ligation of eGFP into the XhoI
 site of QCX (Julius *et al.*, Q vectors, bicistronic retroviral vectors for gene transfer.
Biotechniques, 28:702-08, 2000). Subsequently, full-length FLAG-ATF5 was blunt ligated
 into the BsiWI site of QCX-eGFP, to form the bicistronic Q vector construct (QC-FLAG-
 ATF5-eGFP) for retrovirus production.

30 [00155] The CRE-luciferase reporter plasmid was constructed by annealing synthetic
 oligo 5'-TCGAGTCATGGTAAAAATGACGTCATGGTAATTATCATGGTAAAAAT

GACGTCATGGTAATTATCATGGTAAAAATGACGTCATGGTAATTA-3' to 5'-AGC
TTAATTACCATGACGTCATTTTTACCATGATAATTACCATGACGTCATTTTTACCA
TGATAATTACCATGACGTCATTTTTACCATGAC-3', to form a double-stranded DNA
(Peters *et al.*, ATF-7, a novel bZIP protein, interacts with the PRL-1 protein-tyrosine
5 phosphatase. *J. Biol. Chem.*, 276:13718-26, 2001). The annealed DNA was ligated into the
XhoI and HindIII sites of the GL3 plasmid. VP16-CREB (Columbia University) was
subcloned into the EcoRI and XbaI sites of the pCMS-eGFP vector.

EXAMPLE 4 – ATF5 ANTISERUM

[00156] The CTRGDRKQKKRDQNK peptide, corresponding to ATF5 DNA-binding-
10 domain I (plus an N-terminal cysteine, for conjugation to keyhole limpet hemocyanin), was
used as the antigen for production of rabbit antiserum.

EXAMPLE 5 – WESTERN-BLOT ANALYSIS

[00157] Cultured cells and adult mouse cortex were harvested in Laemmli sample
buffer. The protein concentrations were measured by the Bradford assay (Bio-Rad, Hercules,
15 CA), and cell proteins were resolved by SDS-PAGE on a 12% gel. The separated proteins
were electrophoretically transferred from the gel to Hybond P membrane (Amersham)
(Towbin *et al.*, Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose
sheets: procedure and some applications. *Proc. Natl. Acad. Sci. U S A*, 76:4350-54, 1979).
The membranes were blocked for 1 h in PBS containing 5% milk and 1% BSA, and
20 immunolabeled overnight with ATF5 antipeptide antiserum, at 1:1000, in PBS containing 5%
milk and 1% BSA. For detection, the blots were washed and probed with goat anti-rabbit
HRP-conjugated antibody (Pierce), and then visualized on film using an enhanced
chemiluminescence detection kit (ECL) (Amersham). For the PC12 cell NGF time course, to
normalize for protein loading, the blots were stripped of immunocomplexes, as described by
25 Amersham, and reprobed with ERK1 C-16 antibody (Santa Cruz) and goat anti-rabbit HRP-
conjugated antibody, followed by ECL film visualization. Densitometry was carried out with
NIH Image 1.62 software.

EXAMPLE 6 – IMMUNOHISTOCHEMISTRY

[00158] For PC12 cells, fluorescence immunohistochemistry was carried out, as
30 previously described (Angelastro *et al.*, Characterization of a novel isoform of caspase-9 that

inhibits apoptosis. *J. Biol. Chem.*, 276:12190-200, 2001). For dissociated telencephalic cultures, the cells were fixed with 4% paraformaldehyde and 2% sucrose in PBS, for 15 min. After 3 washes in PBS, the cells were blocked in 10% non-immune goat serum and 0.3% Triton X-100 for 1 h. The cultures were immunolabeled separately with the following combinations: (1) rabbit anti-GFP (1:1000 dilution; Clontech) and mouse anti-nestin (1:500; rat-401 from the DSBH antibody collection, University of Iowa); (2) rabbit anti-GFP (1:1000 dilution) and mouse TUJ1 (1:2000 dilution; Covance); (3) mouse GFP (1:500; Sigma) and rabbit anti-neurofilament 160 kDa (1:200; Columbia University); or (4) mouse GFP (1:500) and rabbit anti-GFAP (1:500; Dako) antibody, in 10% non-immune goat serum and 0.3% Triton X100, for 1 h, followed by secondary labeling with goat FITC-conjugated anti-rabbit or rhodamine-conjugated anti-mouse antibodies (Alexa) at 1:5000.

[00159] For immunolabeling, embryos were fixed in 4% paraformaldehyde in 0.1 M phosphate buffer, overnight, then cryoprotected in 30% sucrose; they were then frozen in O.C.T. compound (Tissue-TEK). Cryosectioned (14 μ m) embryos were blocked for 1 h in 10% non-immune goat serum and 0.3% Triton X-100. The sections were then incubated with ATF5 antiserum (1:500) and TUJ1 antibody (1:2000), in 2.5% nonimmune goat serum and 0.3% Triton X-100, overnight. The sections were subsequently incubated for 1 h with goat FITC-conjugated anti-rabbit and rhodamine-conjugated anti-mouse antibodies, in 10% non-immune goat serum and 0.3% Triton X-100.

[00160] For adherent neurospheres, cells were fixed with 4% para-formaldehyde in PBS / 2% sucrose for 10 min, at room temperature, and then permeabilized for 5 min with 0.5% Triton X-100 in ice-cold PBS / 2% sucrose. After blocking with 25% goat or bovine serum in PBS, for 20 min, the cultures were incubated with primary antibodies (diluted in 25% serum in PBS for 30 min, at room temperature, followed by 3 washes with PBS), and then incubated with the appropriate secondary goat anti-rabbit or anti-mouse antibodies conjugated with FITC (Alexa Fluor 488, Molecular Probes, A 11001) or Texas Red-X (Molecular Probes, T 6391) for 30 min, at room temperature, at 1:200. The cultures were then incubated with the second set of primary and secondary antibodies, as above. Immunochemical reagents were anti-AC133/2 antibody (Miltenyl Biotech), anti-neurofilament 160 (clone NN18, Sigma), anti-beta tubulin isotype III (clone SDL.3D10,

Sigma), and goat anti-tau antiserum (clone C-17, Santa Cruz), all diluted according to the manufacturers' recommendations.

[00161] Confocal microscopy was carried out on either a Zeiss LSM 410 confocal laser scanning microscope (neural brain sections) or on a Bio-Rad Confocal Microscope System 1024ES (neurosphere cultures). Images were obtained under conditions that were identical for both fluorochromes. Confocal images of XY and YZ planes confirmed co-localization in brain sections.

EXAMPLE 7 – IN SITU HYBRIDIZATION

[00162] Non-radioactive *in situ* hybridization of sections was carried out as previously described (Mendelsohn *et al.*, Stromal cells mediate retinoid-dependent functions essential for renal development. *Development*, 126:1139-48, 1999). The antisense ATF5 probe was synthesized using T3 RNA polymerase, and the pCMS-*eGFP-ATF5* construct was digested with NheI as the template. The corresponding sense probe was synthesized using T7 RNA polymerase, and the pCMS-*eGFP-ATF5* construct was digested with NotI as the template.

EXAMPLE 8 – TRANSIENT TRANSFECTIONS

[00163] For PC12 cells, transfection was carried out using 0.5 µg of plasmid / well and 6 µl / well of LipofectAMINE 2000, for 9 h. Thereafter, the cells were re-fed with fresh culture medium, and then handled as described. For telencephalic cells, transfection was performed with 2.0 µg of plasmid / well and 2 µl / well of LipofectAMINE 2000 for 7 h followed by an exchange of medium. For transfection of ATF5 siRNA (AAN19; AAG UCA GCU GCU CUC AGG UAC), 6.67 µg / well of pCMS-EGFP vector were mixed with 80 pmol / well of siRNA in 100 µl of DMEM medium. An equal amount of DMEM medium, premixed with 1 µl of LipofectAMINE 2000 / well, was added to, and mixed with, the vector and siRNA. After 30 min, the final mixture was added to 1/6 the volume containing the cells, and the cells were re-fed with fresh culture medium after 7 h of transfection. For the control, telencephalic cells were transfected with pCMS-EGFP vector alone.

EXAMPLE 9 – RETROVIRUS PRODUCTION AND INFECTION OF TELENCEPHALIC CELLS

[00164] Nonreplicating retrovirus was made by transfecting subconfluent GP2 293 cells (grown in DMEM plus 10% FBS) with 5 μ g of QCX-eGFP or pLeGFP, and 5 μ g of pVSV-G, for production of empty eGFP retrovirus (as described by Clontech). Similarly, GP2 293 cells were transfected with 5 μ g of QC-FLAG-ATF5-eGFP or pLeGFP-FLAG-NTAzip-ATF5, and 5 μ g of pVSV-G, to make the bicistronic FLAG-ATF5-eGFP or fusion eGFP-FLAG-NTAzip-ATF5 retroviruses, respectively. After 48 h, medium was collected, and the virus was concentrated by centrifugation at 50,000 xg, at 4°C. The final titer was approximately 1×10^6 virus particles per ml. The telencephalic cells were infected with 5-10 μ l of retrovirus, one day after plating, and the cells were fixed 7 days after infection.

10 EXAMPLE 10 – SCORING OF NEURONAL DIFFERENTIATION

[00165] Transfected cells were detected by positive immunostaining for eGFP. Co-staining with anti-FLAG established that the GFP-positive PC12 cells also expressed ATF5 constructs. NGF-treated PC12 cells (transfected unless otherwise noted) were scored for processes of length greater than two cell diameters (about 20 μ m) (Greene *et al.*, Culture and Experimental Use of the PC12 Rat Pheochromocytoma Cell Line. In: *Culturing Nerve Cells*, 2nd ed., Goslin, G.K., ed. (Cambridge, MA: The MIT Press, 1998) pp. 161-87. Transfected telencephalic neurons were scored for the presence of processes with lengths greater than two cell diameters (about 20 μ m) and for co-staining with TUJ1, nestin, or NF-M antisera antibodies.

20 EXAMPLE 11 – CRE-LUCIFERASE REPORTER ASSAY

[00166] PC12 cells were co-transfected with 1 μ g of pCMS-eGFP (empty, or containing FLAG-tagged-ATF5 or FLAG-tagged NTAzip-ATF5) and with 0.2 μ g of pGl3-CRE-luciferase reporter and 1 μ g of LacZ plasmid per well; cells were then transfected with 2 μ l / well of LipofectAMINE 2000 24 h prior to harvesting. The cells were treated with NGF for a total of 1 h to 3 days. Luciferase levels were assayed using the Promega Luciferase System with Reporter Lysis Buffer, as described by the manufacturer. The level of LacZ activity was measured, as previously described (Sambrook *et al.*, Molecular Cloning. In: *A Laboratory Manual*, 2nd ed. (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 1989) pp. 16-66.

30 EXAMPLE 12 – STATISTICAL ANALYSES

[00167] Multiple comparisons among the data from different plasmid transfections and retrovirus infections were achieved using Tucky's one way ANOVA test. Comparisons for pairs of data were conducted with Student's t-distribution test.

5 [00168] Discussed below are results obtained by the inventors in connection with the experiments of Examples 1-12:

Reciprocal Effects of NGF on ATF5 Protein Expression and Neurite Outgrowth

[00169] The inventors' previous findings revealed that long-term NGF treatment promotes a 25-fold down-regulation of ATF5 transcripts in PC12 cells (Angelastro *et al.*,
10 Identification of diverse nerve growth factor-regulated genes by serial analysis of gene expression (SAGE) profiling. *Proc. Natl. Acad. Sci. U S A*, 97:10424-429, 2000). To determine whether this is reflected at the level of protein expression, the inventors cloned the coding sequence of rat ATF5 (GenBank/EBI accession number AY123225), and an antiserum was raised against a peptide corresponding to a portion of the deduced sequence of
15 the DNA-binding domain. Western immunoblotting with this antiserum detected a single major band in extracts of PC12 cells (FIG. 1A), and in extracts of HEK-293 cells, primary human neuroblastoma, and mouse brain (data not shown), with an apparent molecular mass of 20-22 kDa. The nucleotide sequence of rat ATF5 indicates two potential in-frame Kozak start sites, and the apparent molecular mass of 20-22 kDa indicates preferential use of the
20 second.

[00170] A time course of ATF5 protein expression in PC12 cells, in response to NGF treatment, revealed a drop in levels by 1 day, and a progressive fall thereafter, with relatively little detectable expression by day 10 (FIGs. 1A and 1B). Quantification of neurite outgrowth in the same sets of cultures revealed a reciprocal relationship with ATF5 expression (FIG.
25 1B).

Exogenous ATF5 Represses NGF-Promoted Neurite Outgrowth, While a Dominant-Negative ATF5 Accelerates Initial Neuritogenesis

[00171] The inverse behaviors of ATF5 expression and neurite outgrowth suggested a possible causal relationship. To test this, FLAG-tagged ATF5 was subcloned into the pCMS-eGFP vector, and transfected into PC12 cells. Two days later, NGF was added, and the
30 transfected cells (expressing eGFP and tagged ATF5) were assessed over time for the

appearance of neurites. In contrast to cells transfected with empty vector, those expressing exogenous ATF5 showed markedly repressed genesis of neurites over a 5-day time course (FIGs. 2A and 2B).

[00172] To assess the possibility that exogenous ATF5 might act, at least in part, by non-physiologically sequestering and "squenching" the actions of binding partners, the inventors also prepared a construct encoding an N-terminally-truncated form of FLAG-tagged ATF5, possessing an enhanced b-Zip domain (NTAzip-ATF5). This was achieved by deleting the N-terminal acidic activation domain, and replacing the DNA-binding domain with an amphipathic acidic α -helical sequence containing leucine repeats at each seventh residue.

[00173] Without activation and DNA-binding domains, NTAzip-ATF5 does not interact with DNA or directly affect gene transcription. However, because this protein includes the intact ATF5 leucine zipper, it retains specific interactions with endogenous ATF5 and with heterologous binding partners. In addition, the Azip amphipathic acidic α -helical domain should tightly associate with the basic DNA-interaction domains of ATF5-binding partners, thereby blocking their functions (Vinson *et al.*, Dimerization specificity of the leucine zipper-containing bZIP motif on DNA binding: prediction and rational design. *Genes Dev.*, 7:1047-58, 1993; Krylov *et al.*, Extending dimerization interfaces: the bZIP basic region can form a coiled coil. *EMBO J.*, 14:5329-37, 1995; Moitra *et al.*, Life without white fat: a transgenic mouse. *Genes Dev.*, 12:3168-81, 1998; Moll *et al.*, Attractive interhelical electrostatic interactions in the proline- and acidic-rich region (PAR) leucine zipper subfamily preclude heterodimerization with other basic leucine zipper subfamilies. *J. Biol. Chem.*, 275:34826-832, 2000). Thus, if exogenous ATF5 acts by non-specific squenching, rather than by binding to DNA, NTAzip-ATF5 should have a similar effect. However, in contrast to ATF5, NTAzip-ATF5 did not block NGF-promoted neurite outgrowth (FIG. 2B), thus ruling out a non-specific action of the former.

[00174] In addition to serving as a control for non-specific squenching, NTAzip-ATF5 acts as a dominant-negative for ATF5, thereby permitting evaluation of the consequences of ATF5 loss-of-function. In the absence of NGF, transfected NTAzip-ATF5 did not stimulate neurite outgrowth (data not shown). However, cells transfected with NTAzip-ATF5, and then exposed to NGF, showed a significantly faster (two-fold) initial appearance of neurites,

as compared with controls (FIG. 2C). This reinforces the notion that a physiologic function of ATF5 is suppression of neurite outgrowth, and that its down-regulation is required for this process to occur. After the first 1-2 days of NGF treatment, the effect of NTazip-ATF5 is much less apparent, presumably due to down-regulation of endogenous ATF5.

5 ATF5 is Highly Expressed in Ventricular Zones of Developing Brain

[00175] The suppression of neurite outgrowth by ATF5 in PC12 cells, and the potential suitability of this system for modeling the transition of neural progenitor cells to differentiated post-mitotic neurons, led the inventors to examine expression of ATF5 in the developing nervous system. *In situ* hybridization revealed specific expression of ATF5 transcripts in E12-15 rat neural nasal epithelium (see, also, Hansen *et al.*, Mouse Atf5: molecular cloning of two novel mRNAs, genomic organization, and odorant sensory neuron localization. *Genomics*, 80:344-50, 2002), dorsal root and trigeminal ganglia, and brain (FIG. 3 and data not shown). The only signal of comparable strength detected outside the nervous system at these stages was in liver (data not shown). Within E12-15 rat brain, expression was highest in the ventricular zone (VZ) of the neural epithelium adjacent to the lateral ventricles and the fourth ventricle – sites of intense proliferation of neural cell precursors – and was decreased in overlying structures containing migrating and post-mitotic neurons (FIG. 3A, panels a and b).

[00176] In view of the pattern of ATF5 transcripts in developing brain, the inventors next examined ATF5 protein expression in developing brain, using immunohistochemistry. ATF5 protein was strongly expressed in the VZ of E12 and E14 telencephalon, and fell to undetectable levels toward the surface of the developing cortex (FIG. 3A, panels c-f; FIG. 3B). Double staining with the TUJ1 antibody that recognizes tubulin β III, a marker for post-mitotic neurons (Lee *et al.*, Posttranslational modification of class III beta-tubulin. *Proc. Natl. Acad. Sci. U S A*, 87:7195-99, 1990), showed a converse pattern of staining (FIGs. 3A and 3B), indicating that ATF5 is highly expressed in proliferating neural progenitor cells and undetectable in differentiated neurons. A comparable pattern was also observed in E14 rat embryo telencephalon, at higher magnification, using confocal microscopy (FIG. 3B). At E17, ATF5 expression remained largely confined to the VZ, in contrast to the large expansion of TUJ1-positive staining in the cortical area (FIGs. 4A-4F).

ATF5 is a Marker For Neural Stem/Progenitor Cells, but not for Mature Neurons in Clonal Neural Progenitor Cell Cultures

[00177] The above findings indicate that ATF5 is highly expressed in proliferating PC12 cells and in VZ progenitor cells, but not in post-mitotic neurons. To further examine the correlation between ATF5 expression and neuronal differentiation, the inventors prepared cultures of neural progenitor cells from the neurogenic subventricular zone or hippocampal dentate gyrus of newborn mouse brain. Clones derived from single-cell suspensions were expanded and cultured as neurospheres, under non-adherent conditions, in the presence of EGF, bFGF, and insulin, and then plated onto poly-L-ornithine and laminin, with 10% fetal bovine serum, to trigger substrate attachment and neurogenesis (Kukekov *et al.*, Multipotent stem/progenitor cells with similar properties arise from two neurogenic regions of adult human brain. *Exp. Neurol.*, 156:333-44, 1999; Laywell *et al.*, Identification of a multipotent astrocytic stem cell in the immature and adult mouse brain. *Proc. Natl. Acad. Sci. U S A*, 97:13883-888, 2000). Cells at the centers of the cultured neurospheres proliferate as stem/progenitor cells, while those that migrate to the culture periphery differentiate into neurons and glia (FIG. 5E).

[00178] ATF5 expression was very high at the 3-dimensional core of the cultures. Co-staining with antibodies to the AC133 antigen, a marker for hematopoietic and neural stem cells (Yin *et al.*, AC133, a novel marker for human hematopoietic stem and progenitor cells. *Blood*, 90:5002-12, 1997; Uchida *et al.*, Direct isolation of human central nervous system stem cells. *Proc. Natl. Acad. Sci. U S A*, 97:14720-25, 2000; Bhatia, AC133 expression in human stem cells. *Leukemia*, 15:1685-88, 2001; Yu *et al.*, AC133-2, a novel isoform of human AC133 stem cell antigen. *J. Biol. Chem.*, 277:20711-716, 2002), revealed extensive co-expression with ATF5 in this region (FIG. 5A). AC133 antigen localization appeared to be largely at the cell surface and plasma membrane, while ATF5 appeared to be mainly localized to nuclei. ATF5 was also extensively expressed in cells positive for nestin (FIG. 5B), an intermediate filament expressed by neuroectodermal progenitors (Lendahl *et al.*, CNS stem cells express a new class of intermediate filament protein. *Cell*, 60:585-95, 1990).

[00179] Co-localization experiments were also carried out with ATF5 and neuronal markers. The 160-kDa neurofilament protein, NF-M, was detected in cells outgrowing towards the culture periphery. A sub-population of such cells, which generally appeared to have short, neurite-like processes, co-stained for nuclear ATF5 (FIG. 5C). For such cells,

staining of ATF5 and NF-M appeared to be of relatively low intensity, indicating that these were immature neuronal cells in transition with rising levels of NF-M expression and falling levels of ATF5. Another population of cells, with more advanced neuronal morphology, strongly stained for NF-M, but was negative for expression of ATF5 (FIG. 5D). Finally, co-
 5 staining with antiserum, for the neuronal marker, tau (Takemura *et al.*, In situ localization of tau mRNA in developing rat brain. *Neuroscience*, 44:393-07, 1991), revealed a set of tau-positive cells, at the periphery of the cultures, with clear neuronal morphology (FIGs. 5E and 5F). Unlike the progenitor cells in the centers of the cultures, that were positive for ATF5 expression and negative for tau, the tau-positive cells in the periphery did not co-stain for
 10 ATF5. Taken together, these observations indicate that ATF5 is expressed in neural stem (AC133+) and progenitor (nestin+) cells, including those committed to the neuronal lineage, and are down-regulated in differentiated, post-mitotic neurons (tau+).

ATF5 Represses, but Dominant-Negative ATF5 and ATF5 Small Interfering RNA Accelerate, Neuronal Differentiation of Neural Progenitor Cells

15 [00180] The above-described expression pattern of ATF5 raised the possibility that the presence of this protein, as in PC12 cells, may block proliferating neural progenitor cells from undergoing neuronal differentiation. To assess this, rat E14 telencephalic cell cultures containing a mixture of proliferating progenitor cells and post-mitotic neurons, and a small number of glial cells (Ghosh and Greenberg, Distinct roles for bFGF and NT-3 in the
 20 regulation of cortical neurogenesis. *Neuron*, 15:89-03, 1995), were transfected with pCMS-eGFP containing either no insert, FLAG-ATF5, or FLAG-NTAzip-ATF5. Transfected cells (identifiable by eGFP expression) were scored 3 days later for neuronal morphology and expression of nestin and tubulin β III (FIG. 6A). In contrast with the cells transfected with empty vector, few cells transfected with ATF5 exhibited neuronal morphology.

25 [00181] In addition, ATF5 greatly repressed expression of the neuronal marker, tubulin β III. On the other hand, ATF5 significantly increased the proportion of cells expressing nestin, a marker for neural progenitor cells. NTAzip-ATF5 did not mimic ATF5, ruling out a potential non-physiological squelching action of ATF5, as in the case of PC12 cells. In comparison with control transfectants, somewhat fewer cells transfected with NTAzip-ATF5
 30 expressed nestin, although a greater number tended to express neuronal markers.

[00182] To ensure initial expression only in proliferating cells of the inventors' telencephalic cell cultures, and to permit transgene delivery at an early point after establishment of the cultures (which was technically unfeasible with the inventors' transfection conditions), the inventors constructed, and infected the cells at 1 day *in vitro* with, retroviral vectors expressing either eGFP, eGFP-FLAG-NTAzip-ATF5, or FLAG-ATF5 and eGFP. In this paradigm, ATF5 once again suppressed neurite outgrowth and expression of neuronal markers (NF-M and TUJ1), and led to an increase in proportion of nestin-positive cells at either 7 (FIG. 6B) or 4 (FIG. 6C) days after infection. Moreover, loss of function of endogenous ATF5, promoted by NTAzip-ATF5, significantly enhanced the genesis of neurite-bearing, TUJ1-positive cells in cultures assessed at 3 (data not shown) and 4 (FIG. 6C) days following viral exposure. The double-negative construct also promoted a fall in nestin positive cells, which presumably reflected the increase in neuronal differentiation. The increase in TUJ1-positive cells was greater than can be accounted for by the fall of nestin-positive cells, indicating either that the antibody the inventors employed led to an underestimation of the numbers of nestin-positive progenitor cells in the cultures, or that at least some neurons were generated from a population of nestin-expressing progenitors.

[00183] To corroborate the inventors' findings that NTAzip-ATF5 accelerates neurogenesis by specifically interfering with the function of endogenous ATF5, rather than through non-specific actions, the inventors employed small interfering RNA (siRNA) to selectively down-regulate endogenous ATF5. After 3 days *in vitro*, E14 telencephalic cells were transfected with GFP or with GFP plus ATF5 siRNA. On the fourth day after transfection with the siRNA, the proportion of transfected cells with detectable endogenous ATF5 fell by 96%, as compared with controls (FIG. 6D). Significantly, the reduction of endogenous ATF5 resulted in a 3.4-fold increase in neurogenesis, as judged by the appearance of TUJ1 staining (FIG. 6D) and neurite outgrowth (data not shown). In contrast, an irrelevant siRNA synthesized to target down-regulation of the protein, POSH, had no effect on development of neuronal markers or processes. Taken together, these findings support a model in which ATF5 suppresses the transition between neural progenitor cells and post-mitotic neurons, and in which loss of, or interference with, ATF5 function accelerates neuronal differentiation.

[00184] The limited degree of neuronal differentiation in the telencephalic cultures appears to occur in response to endogenous factors. To determine whether ATF5 can also regulate CNS neuronal differentiation promoted by a defined trophic agent, the inventors tested the effects of exogenous ATF5 and NTAzip-ATF5 in the presence and absence of

5 NT3, a neurotrophin previously reported to drive telencephalic progenitor cell differentiation into neurons (Ghosh and Greenberg, Distinct roles for bFGF and NT-3 in the regulation of cortical neurogenesis. *Neuron*, 15:89-03, 1995). As shown in FIG. 6E, NT3 nearly tripled the level of neurogenesis in the cultures, and ATF5 suppressed this by 5- to 6-fold.

Contrastingly, NTAzip-ATF5 had no significant effect on neurogenesis in the presence of

10 NT3 – unlike its marked promotion of neuronal differentiation in the absence of NT3. The latter observation would suggest that neuronal differentiation in the cultures is maximally stimulated by NT3, and cannot be further promoted by interfering with endogenous ATF5 activity. Furthermore, it appears that NT3 leads to down-regulation of endogenous ATF5, as none of the neurons formed in its presence exhibited detectable ATF5 immunostaining (data

15 not shown). In conclusion, these findings indicate that, as in the case of NGF, NT3 promotes neurogenesis by a mechanism that can be suppressed by exogenous ATF5, and which includes loss of endogenous ATF5 expression.

Inhibition of Neurite Outgrowth by ATF5 Involves Repression of CRE Transactivation

[00185] The work of Peters *et al.* (ATF-7, a novel bZIP protein, interacts with the

20 PRL-1 protein-tyrosine phosphatase. *J. Biol. Chem.*, 276:13718-26, 2001) has established that ATF5 homodimers specifically bind to CRE elements, and that there is evidence that CRE plays an important role in neuronal differentiation and maintenance (Finkbeiner *et al.*, CREB: a major mediator of neuronal neurotrophin responses. *Neuron*, 19:1031-47, 1997). Hence, the inventors next determined whether ATF5 regulates CRE activity in neuronal cells, and

25 whether this action plays a role in ATF5-mediated suppression of neuronal differentiation.

[00186] The inventors also wished to determine whether the presence of NGF would affect the capacity of ATF5 to regulate CRE activity. Accordingly, PC12 cells were co-transfected with a CRE-luciferase reporter construct, a lacZ expression construct (for normalization of transfection efficiency), and pCMS-eGFP containing either no insert,

30 FLAG-ATF5, or FLAG-NTAzip-ATF5. One day later, the cells were harvested and assessed for reporter activity. A portion of the cultures were treated with NGF for 2 days prior to, and

during, the 24 h after transfection (3-day NGF treatment); others were either unexposed to NGF, or exposed to the factor at the time of transfection (1-day NGF treatment) or during the last hour before harvesting (1-h NGF treatment).

[00187] Without NGF treatment, or after 1 h of NGF treatment, there was relatively
5 little constitutive CRE transactivation. The effect of exogenous ATF5 was somewhat
variable at this time, with suppression of activity in some experiments and not others (FIGs.
7A and 7B), possibly reflecting cell-culture conditions. At 1 day with NGF, there was a
small (50%), but statistically significant, increase in CRE activity, in comparison with naïve
cells; this was reduced to baseline by exogenous ATF5. At day 3, there was a 10-fold
10 increase in CRE reporter activity, as compared with untreated cells, and this was again
substantially reduced by exogenous ATF5. NTazip-ATF5 did not reduce CRE activity,
thereby making it unlikely that ATF5 interferes with CRE transactivation by non-physiologic
interaction with CRE-regulatory proteins. Moreover, neither ATF5 nor NTazip-ATF5
expression suppressed expression of a SRE reporter (data not shown). In addition to
15 establishing that ATF5 suppresses CRE transactivation in intact neuronal cells, these findings
indicate that NGF elevates basal CRE activity, and that this occurs at a time when
endogenous ATF5 levels have fallen by about 2/3 (FIG. 1).

[00188] If ATF5 suppresses neuronal differentiation by binding to CRE and inhibiting
its transactivation, then one might predict that this action should be reversed, either by a
20 dominant-negative ATF5 protein without DNA binding or activation sites, or by a strong
competitive CRE activator. The former characteristics are fulfilled by NTazip-ATF5, which
should form tight heterodimers with ATF5, but does not bind DNA. In support of the
inventors' hypothesis, co-expression of NTazip-ATF5 blocked inhibition of CRE reporter
activity by ATF5 (FIG. 7B), and reversed ATF5-dependent suppression of NGF-promoted
25 neurite outgrowth (FIG. 7C).

[00189] With respect to a competitive CRE activator, the inventors employed VP16-
CREB, a constitutively-active form of the CRE-binding protein, CREB (Lu *et al.*, The
herpesvirus transactivator VP16 mimics a human basic domain leucine zipper protein, luman,
in its interaction with HCF. *J. Virol.*, 72:6291-97, 1998; Barco *et al.*, Expression of
30 constitutively active CREB protein facilitates the late phase of long-term potentiation by
enhancing synaptic capture. *Cell*, 108:689-03, 2002). Co-transfection of pCMS-eGFP-VP16-

CREB into PC12 cells produced strong transactivation of the CRE reporter (FIG. 7A), and this was essentially unaffected by the additional co-transfection of FLAG-ATF5 (FIG. 7B).

[00190] The inventors next assessed whether driving CRE with VP16-CREB would reverse the actions of ATF5 on neurite outgrowth. Transfection of PC12 cells with pCMS-eGFP-VP16-CREB alone did not elicit neurite outgrowth in the absence of NGF, and, as in the case of FLAG-NTAzip-ATF5, enhanced the initial rate of neuritogenesis in the presence of NGF (FIG. 7C). Significantly, co-transfection of VP16-CREB, along with FLAG-ATF5, reversed the suppression of NGF-stimulated neurite outgrowth that was achieved with ATF5 alone (FIG. 7C). Taken together, these findings further support a model in which CRE transactivation is required for neuronal differentiation, but is reversibly blocked by ATF5.

Regulation of Endogenous ATF5 Protein in PC12 Cells and Neural Progenitor Cells

[00191] In consonance with their past observations of ATF5 transcripts, the inventors found that ATF5 protein is expressed in PC12 cells, and drops to nearly undetectable levels during NGF-promoted neuronal differentiation. Similarly, both ATF5 transcripts and protein are highly expressed in neural progenitor cells, and absent from post-mitotic neurons. The observed decrease in ATF5 protein expression most likely reflects the down-regulation of ATF5 transcripts. ATF5 has been reported to be a substrate for ubiquitin-conjugating enzymes, including Cdc34 (Pati *et al.*, Human Cdc34 and Rad6B ubiquitin-conjugating enzymes target repressors of cyclic AMP-induced transcription for proteolysis. *Mol. Cell Biol.*, 19:5001-13, 1999); thus, it is likely to have a relatively rapid turnover that would produce efficient loss of expression following transcriptional down-regulation.

[00192] Western immunoblotting permitted the inventors to deduce the major cellular form of ATF5 protein. The ATF5 cDNA sequence predicts two potential in-frame methionine start sites that would lead to proteins of approximately 30 and 20 kDa. The inventors' observation that the major form of ATF5 in cells has an apparent molecular mass of 20-22 kDa indicates favored utilization of the second site. When a canonical Kozak initiation consensus sequence was included upstream of the first methionine, the larger protein was expressed (data not shown), thereby indicating that the 22-kDa form is not formed by cleavage of a 30-kDa precursor.

ATF5 Represses Neuronal Differentiation of Neural Progenitor Cells

[00193] The down-regulation of ATF5 expression by NGF in PC12 cells, the progressive loss of ATF5 expression that occurs as cells leave the ventricular zone and enter the developing cortex, and the presence of ATF5 in neural stem and progenitor cells, but not in well-differentiated neurons in neurosphere cultures, suggested to the inventors that this factor may play a causal role in regulating neuronal differentiation. In support of this supposition, exogenous ATF5 suppressed both neurite outgrowth in PC12 cell cultures and differentiation of cultured neural progenitor cells. Conversely, loss of ATF5 function (evoked by NTAzap, an ATF5 dominant-negative) nearly doubled the initial rate of NGF-promoted neuritogenesis by PC12 cells, and significantly enhanced neurogenesis in telencephalic cell cultures. In particular, an ATF5 siRNA that effectively reduced endogenous ATF5 levels also promoted a 3.6-fold enhancement of neurogenesis by cultured telencephalic cells.

[00194] The effect of exogenous ATF5 does not appear to be limited solely to neurite outgrowth, as virally-induced ATF5 expression in proliferating progenitor cells also blocked the appearance of several neuronal markers and led to an increase in numbers of cells that expressed nestin – a marker for neural progenitor cells. The increase in numbers of nestin-positive cells induced by exogenous ATF5 appeared to be greater than could be accounted for merely by simply blocking progenitor-cell differentiation. One possible explanation is that nestin-positive cells expressing exogenous ATF5 continued to proliferate, instead of leaving the cell cycle and differentiating.

[00195] Taken together, the inventors' observations with developing rat brain and neurosphere cultures indicate a scenario in which ATF5 is highly expressed in neural stem cells and neuroprogenitor cells, and suppresses their differentiation. The action of appropriate neurotrophic factors leads to down-regulation of ATF5, thereby permitting differentiation of neural progenitor cells into neurons. Therefore, the inventors' present findings suggest that ATF5 acts in a permissive, rather than instructional, manner, in that it does not appear to play a role in directly specifying cell fate *per se*; rather, it appears to act as a negative suppressor that must be down-regulated to permit the transition of neural progenitor cells to neurons. In this role, ATF5 would function to prevent stem cells and progenitor cells from undergoing terminal differentiation until stimulated by appropriate neurotrophic agents.

[00196] Further support for the notion that ATF5 acts as a negative permissive regulator, rather than as an instructional factor, comes from the inventors' observations with NTAzip-ATF5. This modified form of ATF5 should act as a dominant-negative that prevents interaction of ATF5 with DNA as well as with other potential protein-binding partners. This is borne out by the capacity of NTAzip-ATF5 to reverse the effect of ATF5 on CRE reporter activity. Nevertheless, when expressed in PC12 cells, NTAzip did not promote neurite outgrowth in the absence of NGF. Thus, although ATF5 down-regulation appears to be necessary for neuronal differentiation, loss of ATF5 activity does not appear to be sufficient to promote this process. Factors such as NGF appear to down-regulate negative permissive agents such as ATF5 and to provide instructional information that actively promotes neuronal differentiation. In the CNS neuroprogenitor cultures employed herein, down-regulation and instructional activity were likely to be supplied by endogenously-synthesized and released factors, such as NT3 and BDNF (Ghosh and Greenberg, Distinct roles for bFGF and NT-3 in the regulation of cortical neurogenesis. *Neuron*, 15:89-03, 1995).

[00197] The expression pattern of NGF during embryogenesis makes it unlikely that this factor is a key regulator of ATF5 expression in developing brain. However, many other potential neurotrophic factors are present there that could fulfill a similar role. For instance, BDNF and NT3, and their cognate receptors, TrkB and TrkC, are present in rat ventricular progenitor cells at E13 and E15 (Fukumitsu *et al.*, Simultaneous expression of brain-derived neurotrophic factor and neurotrophin-3 in Cajal-Retzius, subplate and ventricular progenitor cells during early development stages of the rat cerebral cortex. *Neuroscience*, 84:115-27, 1998). BDNF (Ahmed *et al.*, BDNF enhances the differentiation but not the survival of CNS stem cell-derived neuronal precursors. *J. Neurosci.*, 15:5765-78, 1995) and NT3 (Ghosh and Greenberg, Distinct roles for bFGF and NT-3 in the regulation of cortical neurogenesis. *Neuron*, 15:89-03, 1995) promote differentiation of cultured neuronal progenitor cells.

[00198] The inventors' experiments were focused on neuronal differentiation, and did not establish whether ATF5 also affects glial cell differentiation. However, the following evidence suggests that ATF5 may also be a negative regulator of astrocyte differentiation: the localization of ATF5 in brain areas that also give rise to glial progenitor cells; the co-localization of ATF5 with nestin, which is present in progenitor cells for both neurons and glia; and the inventors' preliminary observations that ATF5 co-localizes with GFAP in

neuroprogenitor cell cultures and that exogenous ATF5 suppresses GFAP expression.

Although ATF5 expression negatively correlates with neuronal differentiation, this may not be the case universally for differentiation of other cell types. Peters *et al.* (ATF-7, a novel bZIP protein, interacts with the PRL-1 protein-tyrosine phosphatase. *J. Biol. Chem.*,

5 276:13718-26, 2001) reported that ATF5 transcripts were markedly elevated when human Caco-2 cells reached confluency and spontaneously differentiated into a brush-border-bearing polarized cell layer.

Suppression of Neuronal Differentiation by ATF5 Involves CRE

[00199] Based on reports that ATF5 homodimers bind CRE, but not C/EBP or AP1, sites (Peters *et al.*, ATF-7, a novel bZIP protein, interacts with the PRL-1 protein-tyrosine phosphatase. *J. Biol. Chem.*, 276:13718-726, 2001), and that ATF5 represses cAMP-mediated activation of a CRE reporter in JEG3 cells (Pati *et al.*, Human Cdc34 and Rad6B ubiquitin-conjugating enzymes target repressors of cyclic AMP-induced transcription for proteolysis. *Mol. Cell Biol.*, 19:5001-13, 1999), the inventors examined the effect of ATF5 on the activity of a CRE reporter in PC12 cells. The inventors' findings confirm that ATF5 suppresses cellular CRE transactivation. As discussed above, it is significant that NTAzip-ATF5 did not mimic the suppressive actions of ATF5 on neurite outgrowth and CRE activity; rather, it antagonized these effects, thereby indicating that ATF5 acts by binding to DNA, instead of by non-specific "squenching" of binding partners.

20 [00200] The inventors observed that basal CRE activity substantially increased by 3 days of NGF treatment. One potential cause for this is the concurrent fall in endogenous ATF5 expression, and subsequent loss of ATF5-mediated CRE repression; however, the inventors cannot rule out the possibility that NGF regulates additional proteins that affect CRE activity.

25 [00201] Although NTAzip-ATF5 blocked the inhibitory effects of exogenous ATF5 on its own, it had no, or relatively little, effect on CRE reporter activity. If, as the inventors propose, CRE-dependent gene activation is suppressed by endogenous ATF5, then it might have been anticipated that basal CRE activation would be elevated in response to NTAzip-ATF5. Since this was not the case, this raises the possibility that one or more factors, in addition to ATF5, act to suppress CRE in neural progenitor cells, and that these are also down-regulated during neuronal differentiation.

30

[00202] To assess whether interference with CRE-mediated gene regulation might account for the mechanism by which ATF5 interferes with neuronal differentiation, the inventors co-expressed it with VP16-CREB, a constitutively-active fusion protein that includes the CREB DNA-binding domain and transactivation domain of the HSV VP16 protein. VP16-CREB potentially activated the CRE reporter, and this effect was not blocked by co-expression of ATF5. Significantly, co-expressed VP16-CREB overrode ATF5-mediated inhibition of neurite outgrowth. This finding supports a model in which neuronal differentiation requires CRE-mediated gene activation, and in which such activation is repressed in neural progenitor cells by factors such as ATF5. In this light, it is of interest that PACAP, a potent activator of adenylate cyclase, promotes mitotic exit and neuronal differentiation of cultured cortical neuron precursor cells (Dicicco-Bloom *et al.*, The PACAP ligand/receptor system regulates cerebral cortical neurogenesis. *Ann. N. Y. Acad. Sci.*, 865:274-89, 1998), and that NGF-promoted differentiation of PC12 cells is synergized by cell-permeant cAMP derivatives (Gunning *et al.*, Differential and synergistic actions of nerve growth factor and cyclic AMP in PC12 cells. *J. Cell Biol.*, 89:240-45, 1981).

[00203] In summary, the inventors' findings indicate that both positive and negative regulators govern the transition of neural progenitor cells to neurons. On one hand, ATF5 is highly expressed in neural stem cells and neuroprogenitor cells, and suppresses their neuronal differentiation, apparently by competing for binding to CREs. On the other hand, neuronal differentiation is accompanied by, and appears to require, down-regulation of ATF5 expression. This can be accomplished by neurotrophic factors such as NGF and NT3. Though such down-regulation may be necessary, it is not sufficient to permit neuronal differentiation. The latter also appears to require instructive signals that may be imparted by neurotrophic factors and/or activators of adenylate cyclase.

[00204] While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be appreciated by one skilled in the art, from a reading of the disclosure, that various changes in form and detail can be made without departing from the true scope of the invention in the appended claims.